

For Research Use Only

VPS52 Polyclonal antibody

Catalog Number: 11662-2-AP 2 Publications



Basic Information

Catalog Number: 11662-2-AP	GenBank Accession Number: BC032108	Purification Method: Antigen affinity purification
Size: 400 µg/ml	GeneID (NCBI): 6293	Recommended Dilutions: WB 1:500-1:1000
Source: Rabbit	UNIPROT ID: Q8N1B4	IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate
Isotype: IgG	Full Name: vacuolar protein sorting 52 homolog (S. cerevisiae)	IHC 1:20-1:200
Immunogen Catalog Number: AG2180	Calculated MW: 723 aa, 82 kDa	IF 1:10-1:100
	Observed MW: 82 kDa	

Applications

Tested Applications: IF/ICC, IHC, IP, WB, ELISA	Positive Controls:
Cited Applications: WB, IHC	WB: HepG2 cells, COLO 320 cells, mouse skin tissue
Species Specificity: human, mouse, rat	IP: mouse skin tissue,
Cited Species: human, mouse	IHC: human gliomas tissue,
	IF: HepG2 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

VPS52 is a component of the Golgi-associated retrograde protein (GARP) complex, also called VFT (VPS fifty-three) complex, composed of VPS51, VPS52, VPS53 and VPS54. GARP complex functions in traffic from endosomes the trans-Golgi network (PMID: 15878329). GARP proteins interact with RAB proteins and SNARE proteins. VPS52 may play a role in the development of a multicellular organism through cell-cell interactions (PMID: 23142660).

Notable Publications

Author	Pubmed ID	Journal	Application
Jian Zhang	28791438	J Mol Med (Berl)	IHC, WB
Jian Wei	28658628	Cell Rep	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 11662-2-AP (VPS52 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human gliomas using 11662-2-AP (VPS52 antibody) at dilution of 1:50 (under 10x lens).



IP result of anti-VPS52 (IP:11662-2-AP, 3ug; Detection:11662-2-AP 1:1000) with mouse skin tissue lysate 3200ug.



Immunofluorescent analysis of HepG2 cells using 11662-2-AP (VPS52 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).