For Research Use Only

NFATC2IP Polyclonal antibody

Catalog Number: 11667-1-AP



Purification Method:

WB 1:500-1:3000 IF/ICC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 11667-1-AP BC021551 GeneID (NCBI): Size: 650 ug/ml 84901 Source: **UNIPROT ID:** Rabbit Q8NCF5 Full Name: Isotype:

nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 2 Immunogen Catalog Number: interacting protein

Calculated MW: 138 aa, 15 kDa Observed MW: 50-65 kDa

Applications

Tested Applications: WB, IF/ICC, ELISA Species Specificity:

human

AG2217

Positive Controls:

WB: Jurkat cells, K-562 cells, Raji cells

IF/ICC: HeLa cells,

Background Information

 $NFATC2IP, also \ named \ as \ NIP45, is \ a \ 419 \ amino \ acid \ protein, which \ contains \ one \ ubiquitin-like \ domain. \ TRAF1 \ is$ associated with a fraction of NFATC2IP in the cytoplasm and prevents its translocation to the nucleus. NFATC2IP In T-helper 2 (Th2) cells, regulates the magnitude of NFAT-driven transcription of a specific subset of cytokine genes, including IL3, IL4, IL5 and IL13, but not IL2.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 11667-1-AP (NFATC2IP antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using NFATC2IP antibody (11667-1-AP) at dilution of 1:400 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).