## For Research Use Only

## DYNLT3 Polyclonal antibody

Catalog Number: 11687-1-AP



**Purification Method:** 

WB 1:1000-1:4000 IHC 1:50-1:500

IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number:

11687-1-AP

BC000968

Size:

GeneID (NCBI):

900 µg/ml

6990

Source:

UNIPROT ID:

Rabbit

P51808

Isotype:

GeneID (NCBI):

Full Name:

dynein, light chain, Tctex-type 3

Immunogen Catalog Number: Calculated MW:
AG2299 116 aa, 13 kDa
Observed MW:

13 kDa

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: U-937 cells, HeLa cells IHC: human ovary tumor tissue,

IF/ICC: HepG2 cells,

## **Background Information**

Dynein light chain Tctex-type 3 (DYNLT3) is a member of the cytoplasmic dynamic protein light chain Tctex family. DYNLT3 mainly plays a regulatory role in mitosis and meiosis, as well as chromosome binding. DYNLT3 is expressed in all cells and tissues. The calculated molecular weight of DYNLT3 is 13 kDa.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




U-937 cells were subjected to SDS PAGE followed by western blot with 11687-1-AP (DYNLT3 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 11687-1-AP (DYNLT3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 11687-1-AP (DYNLT3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using DYNLT3 antibody (11687-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).