

For Research Use Only

DDX39A Polyclonal antibody

Catalog Number: 11723-1-AP **5 Publications**



Basic Information

Catalog Number: 11723-1-AP	GenBank Accession Number: BC032128	Purification Method: Antigen affinity purification
Size: 400 µg/ml	GeneID (NCBI): 10212	Recommended Dilutions: WB 1:500-1:1000
Source: Rabbit	UNIPROT ID: O00148	IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate
Isotype: IgG	Full Name: DEAD (Asp-Glu-Ala-Asp) box polypeptide 39	IHC 1:50-1:500
Immunogen Catalog Number: AG2311	Calculated MW: 427 aa, 49 kDa	IF 1:200-1:800
	Observed MW: 49-55 kDa	

Applications

Tested Applications: IF/ICC, IHC, IP, WB, ELISA	Positive Controls: WB : HEK-293 cells, HeLa cells, HepG2 cells, Jurkat cells
Cited Applications: WB, IP, IF, IHC	IP : HEK-293 cells,
Species Specificity: human, mouse, rat	IHC : mouse kidney tissue,
Cited Species: human, chicken, mouse	IF : HepG2 cells,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

DDX39A, also named the BAT1 protein, contains the nine conserved motifs that characterize the DEAD-box family of RNA-binding proteins. The family includes proteins found in all eukaryotic cell types, with considerable divergence in the sequences lying between the conserved motifs. Some of the motifs were known before the definition of the family and are responsible for binding to mRNA or ATP, or possess ATPase activity. Phylogenetic analyses have grouped BAT1 with the defining member of the DEAD-box family, eIF-4. This is a translation initiation factor required for the dissociation of stem/loop structures in mRNA at the ribosomes. This antibody may cross-react with DDX39B.

Notable Publications

Author	Pubmed ID	Journal	Application
Lingcai Zhao	36084138	PLoS Pathog	IF, WB
J J David Ho	34644561	Cell Rep	WB
Yewei Bao	34421357	Int J Biol Sci	WB, IHC

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11723-1-AP (DDX39A antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 11723-1-AP (DDX39A antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using DDX39A antibody (11723-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



IP result of anti-DDX39A (IP:11723-1-AP, 4ug; Detection:11723-1-AP 1:1000) with HEK-293 cells lysate 3000ug.