For Research Use Only

VWF, VWFpp Polyclonal antibody

Catalog Number: 11778-1-AP 107 Publications




Basic Information	Catalog Number: 11778-1-AP	GenBank Accession Number:	Purification Method: Antigen affinity purification
	Size:	GeneID (NCBI):	Recommended Dilutions:
	550 µg/ml	7450	WB 1:1000-1:4000
	Source: Rabbit	UNIPROT ID: P04275	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
	Isotype: IgG	Full Name: von Willebrand factor	
	Immunogen Catalog Number: AG2386	Calculated MW: 309 kDa	
		Observed MW: 309-320 kDa, 75-100 kDa	
Applications	Tested Applications:	Positive	Controls:
	WB, IP, ELISA		VEC cells, Jurkat cells, human placenta tissue
	Cited Applications: WB		spleen tissue, rat spleen tissue
	Species Specificity: human, mouse, rat	IP : mou	se spleen tissue,
	Cited Species:		
	human, mouse, rat, pig, monkey		
Background Information	human, mouse, rat, pig, monkey Von Willebrand factor (vWF) is a platelet adhesion and carries fact common inherited bleeding disor 22-aa signal peptide, a 741-aa la 2,050 aa making up the mature v targeting to storage granules. Thi	tor VIII (FVIII). Deficiency of vWF res rder. The pre-pro-vWF is comprised o rge propeptide (vWF propeptide, vW WF. The von Willebrand antigen 2 is s antibody raised against the N-term	ults in von Willebrand disease (vWD), a f 2,813 amino acids (aa) which encompass a Fpp, also called von Willebrand antigen 2) an
	human, mouse, rat, pig, monkey Von Willebrand factor (vWF) is a platelet adhesion and carries fact common inherited bleeding disor 22-aa signal peptide, a 741-aa la 2,050 aa making up the mature v targeting to storage granules. Thi Willebrand antigen 2 (75-100 kDa	tor VIII (FVIII). Deficiency of vWF res rder. The pre-pro-vWF is comprised o rge propeptide (vWF propeptide, vW WF. The von Willebrand antigen 2 is s antibody raised against the N-term a) and pro-vWF (309-320 kDa). (PMID	f 2,813 amino acids (aa) which encompass a Fpp, also called von Willebrand antigen 2) an required for multimerization of vWF and for i inal region of pre-pro-vWF recognizes von 9759493; 12067899; 22452980; 8874191)
Background Information Notable Publications	human, mouse, rat, pig, monkey Von Willebrand factor (vWF) is a platelet adhesion and carries fact common inherited bleeding disor 22-aa signal peptide, a 741-aa la 2,050 aa making up the mature v targeting to storage granules. Thi Willebrand antigen 2 (75-100 kDa	tor VIII (FVIII). Deficiency of vWF rest rder. The pre-pro-vWF is comprised of rge propeptide (vWF propeptide, vW WF. The von Willebrand antigen 2 is s antibody raised against the N-term a) and pro-vWF (309-320 kDa). (PMIC Pubmed ID Journal	ults in von Willebrand disease (vWD), a f 2,813 amino acids (aa) which encompass a Fpp, also called von Willebrand antigen 2) an required for multimerization of vWF and for i sinal region of pre-pro-vWF recognizes von b: 9759493; 12067899; 22452980; 8874191) Application
	human, mouse, rat, pig, monkey Von Willebrand factor (vWF) is a platelet adhesion and carries fact common inherited bleeding disor 22-aa signal peptide, a 741-aa la 2,050 aa making up the mature vi targeting to storage granules. Thi Willebrand antigen 2 (75-100 kDa Author Yixin Ren	tor VIII (FVIII). Deficiency of vWF res rder. The pre-pro-vWF is comprised o rge propeptide (vWF propeptide, vW WF. The von Willebrand antigen 2 is s antibody raised against the N-term a) and pro-vWF (309-320 kDa). (PMID	ults in von Willebrand disease (vWD), a f 2,813 amino acids (aa) which encompass a Fpp, also called von Willebrand antigen 2) ar required for multimerization of vWF and for innal region of pre-pro-vWF recognizes von b: 9759493; 12067899; 22452980; 8874191) Application


Aliquoting is unnecessary for -20 $^{\circ}$ C storage

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





IP result of anti-VWF, VWFpp (IP:11778-1-AP, 4ug; Detection:11778-1-AP 1:300) with mouse spleen tissue lysate 5200ug. Various lysates were subjected to SDS PAGE followed by western blot with 11778-1-AP (VWF, VWFpp antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.