For Research Use Only

KYNU Polyclonal antibody Catalog Number:11796-1-AP Featured Product

Featured Product






Basic Information	Catalog Number: 11796-1-AP	GenBank Accession Number: BC000879		Purification Method: Antigen affinity purification	
	Size: GenelD (NCBI):			Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500 IF/ICC 1:250-1:1000	
	500 ug/ml	8942			
	Source: UNIPROT ID: Rabbit Q16719 Isotype: Full Name:				
	IgG Immunogen Catalog Number:	kynureninase (L-kynurenine hydrolase)			
	AG2365	Calculated MW: 465aa,52 kDa; 3	07aa,35 kDa	Da	
	Observed MW: 45-52 kDa				
Applications	Tested Applications:	Positive Controls:			
	WB, IHC, IF/ICC, ELISA		WB : HeLa cells, A549 cells, HepG2 cells, rat liver tiss		
	Cited Applications: WB, IHC, IF	IHC : human breast cancer tissue,			
	Species Specificity: IF/ICC : HeLa cells, HepG2 cells human, mouse, rat				
	Cited Species: human, mouse, rat				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	KYNU(kynureninase) is also named as L-kynurenine hydrolase and belongs to the kynureninase family. It is a pyridoxal-5'-phosphate-(pyridoxal-P)-depe11dent enzyme that catalyses the cleavage of L-kynurenine and L-3-hydroxykynurenine into anthranilic and 3-hydroxyanthranilic acids, respectively(PMID:8706755). In SDS-PAGE, it can detect a 48 kDa band lower than the preidicted molecular weight of 52 kDa due to limited proteolysis of the enzyme which might occur during handling of the protein(PMID:8706755). This protein is a 95-kD kD homodimer predominantly located in the cytoplasm(PMID:9180257). It has 2 isoforms produced by alternative splicing.				
Notable Publications	Author	Pubmed ID	Journal	Application	
	C Ci	31419330	Clin Exp Dermatol		
	Yingying Song		Virol Sin	WB	
	Bo Wang	31115516	Mol Med Rep	WB	
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	d 50% glycerol pH 7.3	5.		


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data

for 1.5 hours.







HeLa cells were subjected to SDS PAGE followed by western blot with 11796-1-AP (KYNU antibody) at dilution of 1:2000 incubated at room temperature Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 11796-1-AP (KYNU antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 11796-1-AP (KYNU antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 11796-1-AP (KYNU antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using KYNU antibody (11796-1-AP) at dilution of 1:500 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of HepG2 cells, using KYNU antibody 11796-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).