For Research Use Only

OLR1/LOX1 Polyclonal antibody

Catalog Number:11837-1-AP

Featured Product

41 Publications




Basic Information	Catalog Number: 11837-1-AP	GenBank Accession Nu BC022295	mber:	Purification Method: Antigen affinity purification	
	Concentration:	GenelD (NCBI):		Recommended Dilutions: WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	750 ug/ml Source:	4973 UNIPROT ID:			
	Rabbit	P78380		proteinlysate	
	Isotype: IgG Immunogen Catalog Number: AG2437	IHC 1:50-1:500 Full Name: oxidized low density lipoprotein (lectin-like) receptor 1		THC 1:50-1:500	
		Calculated MW: 273 aa, 31 kDa			
		Observed MW: 50-55 kDa			
Applications	Tested Applications:	Positive Controls:			
	WB, IHC, IP, ELISA Cited Applications:	WB : bEnd.3 cells, human liver tissue			
	WB, IHC, IF, CoIP		IP : LO2 cells,		
	Species Specificity:	IHC : human heart tissue, human placenta tissue			
	human, mouse				
	human, mouse Cited Species: human, mouse				
	Cited Species:	atively, antigen			
Background Informatior	Cited Species: human, mouse Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 OLR1/LOX-1 acts as a receptor, in degradation of oxidatively modif predominantly expressed in endo	atively, antigen d with citrate the form of homodimer, that fied low density lipoprotein othelial cells and vascular-ri n containing 273 amino acic Da, which may result from th	(oxLDL) (PMIE ch organs suc ds with a calcu ne glycosylati	h as the placenta, lung, liver, and bra Jlated molecular mass of 31 kDa but	
	Cited Species: human, mouse Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 OLR1/LOX-1 acts as a receptor, in degradation of oxidatively modif predominantly expressed in endo (PMID: 9828121). OLR1 is a protei apparent molecular mass of 50 kl glycosylation sites located at the	atively, antigen d with citrate the form of homodimer, that fied low density lipoprotein othelial cells and vascular-ri n containing 273 amino acic Da, which may result from th C-terminal domain (PMID: 9	(oxLDL) (PMIE ch organs suc ds with a calcu ne glycosylati	D: 9052782, 15695803). ORL1 is h as the placenta, lung, liver, and bra Jlated molecular mass of 31 kDa but on of four potential N-linked	
	Cited Species: human, mouse Note-IHC: suggested antiger tre buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 OLR1/LOX-1 acts as a receptor, in degradation of oxidatively modif predominantly expressed in endo (PMID: 9828121). OLR1 is a protei apparent molecular mass of 50 kl glycosylation sites located at the	atively, antigen d with citrate the form of homodimer, that fied low density lipoprotein othelial cells and vascular-ri n containing 273 amino acic Da, which may result from th	(oxLDL) (PMIE ch organs suc ds with a calcu ne glycosylati 9052782).): 9052782, 15695803). ORL1 is h as the placenta, lung, liver, and bra Jlated molecular mass of 31 kDa but	
Background Informatior	Cited Species: human, mouse Note-IHC: suggested antiger TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 OLR1/LOX-1 acts as a receptor, in degradation of oxidatively modif predominantly expressed in endd (PMID: 9828121). OLR1 is a protei apparent molecular mass of 50 kl glycosylation sites located at the Author P Meng Zhang 2	atively, antigen d with citrate the form of homodimer, that fied low density lipoprotein thelial cells and vascular-ri n containing 273 amino acio Da, which may result from th C-terminal domain (PMID: 9 ubmed ID Journal	(oxLDL) (PMIE ch organs suc ds with a calcu e glycosylati 9052782).	0: 9052782, 15695803). ORL1 is h as the placenta, lung, liver, and bra Jlated molecular mass of 31 kDa but on of four potential N-linked Application	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



bEnd.3 cells were subjected to SDS PAGE followed by western blot with 11837-1-AP (OLR1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



IP result of anti-OLR1 (IP:11837-1-AP, 4ug; Detection:11837-1-AP 1:800) with LO2 cells lysate 1800ug.



Immunohistochemical analysis of paraffin-embedded human heart tissue slide using 11837-1-AP (OLR1 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human heart tissue slide using 11837-1-AP (OLR1 Antibody) at dilution of 1:200 (under 40x lens).

Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 11837-1-AP (OLR1/LOX1 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).