For Research Use Only

Caspase 4/p20/p10 Polyclonal antibody



Catalog Number: 11856-1-AP

Featured Product

12 Publications

UNIPROT ID:

P49662

peptidase

Calculated MW: 377 aa. 43 kDa Observed MW: 43-48 kDa

Basic Information

Catalog Number: 11856-1-AP Size: $600 \mu g/ml$ Source: Rabbit

Full Name: Isotype: caspase 4, apoptosis-related cysteine

Immunogen Catalog Number:

AG2429

Applications

Tested Applications: WB,IF/ICC,IHC,ELISA Cited Applications:

WB,IHC

Species Specificity:

human **Cited Species:** human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

GenBank Accession Number: **Purification Method:**

BC017839 Antigen affinity purification GeneID (NCBI): Recommended Dilutions:

> WB 1:500-1:1000 IHC 1:20-1:200 IF 1:50-1:500

Positive Controls:

WB: SGC-7901 cells, THP-1 cells IHC: human pancreas cancer tissue,

IF: HepG2 cells,

Background Information

Caspase 4 (CASP4), is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspase 4 is able to cleave and activate its own precursor protein, as well as caspase 1 precursor. Overexpression of caspase 4 will induce cell apoptosis.

Notable Publications

Author	Pubmed ID	Journal	Application
Liangmin Fu	36248876	Front Immunol	IHC,WB
Xiao-Lei Xin	31542790	Med Sci Monit	WB
Qingshan Ji	33118609	Inflammation	WB


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




SGC-7901 cells were subjected to SDS PAGE followed by western blot with 11856-1-AP (CASP4 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human pancreas cancer using 11856-1-AP (CASP4 antibody) at dilution of 1:50 (under 25x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 11856-1-AP (Caspase 4/p20/p10 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 11856-1-AP (Caspase 4/p20/p10 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 11856-1-AP (Caspase 4/p20/p10 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.