## For Research Use Only

# DGKE Polyclonal antibody

Catalog Number: 11900-1-AP

3 Publications



**Basic Information** 

Catalog Number:

11900-1-AP

BC022297

Size:

GeneID (NCBI):

850 µg/ml

8526

Source:

UNIPROT ID:

Rabbit

P52429

Isotype:

Full Name:

Antigen affinity purification Recommended Dilutions: WB 1:1000-1:6000 IHC 1:50-1:500 IF/ICC 1:200-1:800

**Purification Method:** 

Isotype:

IgG diacylglycerol kinase, epsilon 64kDa

Immunogen Catalog Number: Calculated MW:

AG2512 64 kDa

Observed MW: 60-64 kDa

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IP, IF Species Specificity: human, mouse, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

WB: HEK-293 cells, Jurkat cells, mouse brain tissue, rat

brain tissue

IHC: human colon cancer tissue,

IF/ICC: A431 cells,

# **Background Information**

DGKE(Diacylglycerol kinase epsilon) is also named as DAGK5 and belongs to the eukaryotic diacylglycerol kinase family. It may be involved mainly in the regeneration of phosphatidylinositol (PI) from diacylglycerol in the PI-cycle during cell signal transduction(PMID:16689942). When expressed in mammalian cells, DGK-epsilon showed specificity for arachidonyl-containing diacylglycerol.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Di Wu	36415997	Obesity (Silver Spring)	WB
Arpad Palfi	27527066	Sci Rep	IHC
Milena Krajewska	38056763	Biochim Biophys Acta Mol Cell Biol Lipids WB,IF,IP	

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 11900-1-AP (DGKE antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed A431 cells using DGKE antibody (11900-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 11900-1-AP (DGKE antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 11900-1-AP (DGKE antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 11900-1-AP (DGKE antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).