For Research Use Only

MAPKSP1 Polyclonal antibody

Catalog Number: 11937-1-AP

1 Publications



Basic Information

Catalog Number: 11937-1-AP

Isotype:

Size: 300 µg/ml Source: Rabbit

IgG

Immunogen Catalog Number: Calculated MW: AG2534 124 aa, 14 kDa Observed MW:

14 kDa

BC026245

8649

GeneID (NCBI):

UNIPROT ID:

Q9UHA4
Full Name:

GenBank Accession Number:

MAPK scaffold protein 1

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000

IHC 1:20-1:200

Applications

Tested Applications: IHC, WB,ELISA Cited Applications:

WB

Species Specificity: human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human liver tissue, LO2 cells, mouse brain tissue, mouse liver tissue

IHC: human prostate cancer tissue, human heart tissue, human liver tissue

Background Information

MAPKSP1, also named as mitogen activated protein kinase kinase 1 interacting protein 1, Map2k1ip1, MEK partner 1, Lamtor3 and MP1. It was first identified as a scaffold protein enhancing the efficiency of the MAPK pathway by facilitating the interaction between MEK1 and ERK1 upon serum stimulation (PMID: 9733512). Later, it has been demonstrated that MAPKSP1 enhances activation of both ERK1 and ERK2 as a response to EGF (PMID: 12479806) and fibronectin (PMID: 15923628). MAPKSP1 is involved in regulating endosomal signaling (PMID: 17178906), and a role in cancer, via MEK and ERK hyperactivation, has been demostrated in pancreatic tumorigenesis (PMID: 24209743). MAPKSP1 has two isoforms with the molecular weight of 13-14 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Yihan Wang	35959437	Front Pharmacol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



human liver tissue were subjected to SDS PAGE followed by western blot with 11937-1-AP (MAPKSP1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer using 11937-1-AP (MAPKSP1 antibody) at dilution of 1:50 (under 10x lens).