

AGR3 Polyclonal antibody

Catalog Number: 11967-1-AP

5 Publications

Basic Information

Catalog Number:

11967-1-AP

Size:

240 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG2577

GenBank Accession Number:

BC058284

GeneID (NCBI):

155465

UNIPROT ID:

Q8TD06

Full Name:

anterior gradient homolog 3 (Xenopus laevis)

Calculated MW:

166 aa, 19 kDa

Observed MW:

19-25 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:50-1:500

IF/ICC 1:20-1:200

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse

Cited Species:

human, chicken

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : SKOV-3 cells, MCF-7 cells

IHC : human breast cancer tissue,

IF/ICC : SKOV-3 cells,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Soma Samanta	29262583	Oncotarget	WB, IHC
Ying Zhu	31682227	Elife	IF
Rui Ye	34177583	Front Pharmacol	IHC, IF, WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 11967-1-AP (AGR3 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0)).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 11967-1-AP (AGR3 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0)).



SKOV-3 cells were subjected to SDS PAGE followed by western blot with 11967-1-AP (AGR3 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of SKOV-3 cells using 11967-1-AP (AGR3 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).