For Research Use Only

## URG4 Polyclonal antibody

Catalog Number:11998-1-AP

Featured Product






Basic Information	Catalog Number: 11998-1-AP	GenBank Accession Number: BC018426		Purification Method: Antigen affinity purification	
	Size: 650 ug/ml	GenelD (NCBI): 55665		Recommended Dilutions: WB 1:500-1:1000	
	Source: Rabbit	UNIPROT ID: Q8TCY9		IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	Isotype: IgG			IHC 1:200-1:1000 IF/ICC 1:50-1:500	
Applications				ontrols:	
	WB, IHC, IF/ICC, IP, ELISA		WB : HEK-2	193 cells,	
	Cited Applications: WB, IHC		IP : HEK-29	IP : HEK-293 cells,	
			nan stomach cancer tissue, human liver cance		
	Cited Species: human			eLa cells,	
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	URG4 (upregulated gene 4), also known as upregulator of cell proliferation (URGCP), is an oncogene which is involved in the development and progression of various tumors. URG4 is upregulated in hepatocellular carcinoma (HCC), ovarian cancer, and other types of cancer. Overexpression of URG4 not only results in tumour progression but also in metastasis and recurrence.				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Yayun Liu	32552851	J Orthop Surg Res	IHC,WB	
	Yuanxi Leng	37832904	Bone	WB	
Storage	Storage: Store at -20°C. Stable for one ye Storage Buffer: PBS with 0.02% sodium azide an Aliquoting is unnecessary for -2	nd 50% glycerol pH	7.3.		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data







HEK-293 cells were subjected to SDS PAGE followed by western blot with 11998-1-AP (URG4 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours. IP result of anti-URG4 (IP:11998-1-AP, 4ug; Detection:11998-1-AP 1:300) with HEK-293 cells lysate 880ug.



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 11998-1-AP (URG4 antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of URG4 antibody (11998-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-URG4 transfected HEK-293 cells.



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using URG4 antibody (11998-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).