For Research Use Only

NDRG2 Polyclonal antibody

Catalog Number: 12015-1-AP

Featured Product

2 Publications



Basic Information

Catalog Number:

12015-1-AP

BC010458

Concentration:

240 ug/ml

57447

Source:

Rabbit

Q9UN36

Isotype:

GenBank Accession Number:

BC010458

GeneID (NCBI):

57447

UNIPROT ID:

Q9UN36

Full Name:

IgG NDRG family member 2

Immunogen Catalog Number:Calculated MW:AG2629371 aa, 39 kDa

Observed MW: 41-43 kDa Purification Method: Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

Applications

Tested Applications: WB, IHC, IP, ELISA Cited Applications: WB, IHC

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: mouse brain tissue, HepG2 cells, rat brain tissue

IP: rat brain tissue,
IHC: mouse brain tissue,

Background Information

NDRG2, or N-myc downstream-regulated gene 2, is a tumor suppressor gene that plays a significant role in cancer development and progression. It is associated with tumor incidence, progression, and metastasis, and is known to regulate tumor-associated genes. NDRG2 is part of the NDRG family, which includes NDRG1, NDRG2, NDRG3, and NDRG4. These proteins are characterized by an esterase/lipase/thioesterase active site serine and an α / β hydrolase fold of approximately 220 amino acids, although they do not exhibit enzyme activity.

Notable Publications

Author	Pubmed ID	Journal	Application
Fenhong Kang	32345304	J Ovarian Res	WB
Zhao Guang	37033917	Front Immunol	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 12015-1-AP (NDRG2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 12015-1-AP (NDRG2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse brain tissue were subjected to SDS PAGE followed by western blot with 12015-1-AP (NDRG2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



WB result of NDRG2 antibody (12015-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-NDRG2 transfected HepG2 cells.



IP result of anti-NDRG2 (IP:12015-1-AP, 4ug; Detection:12015-1-AP 1:1000) with rat brain tissue lysate 4000ug.