## For Research Use Only

# RTN3 Polyclonal antibody

Catalog Number: 12055-2-AP

19 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 12055-2-AP BC011394 GeneID (NCBI): Size: 450 μg/ml 10313 **UNIPROT ID:** Source: Rabbit 095197 Full Name: Isotype: reticulon 3 Calculated MW: Immunogen Catalog Number: AG2685 113 kDa

> Observed MW: 24 kDa, 100-113 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

IHC 1:50-1:500

WB 1:1000-1:8000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

**Applications** 

Tested Applications: IHC, IP, WB, ELISA Cited Applications: WB, IP, IF

Species Specificity: human Cited Species:

human, rat, mouse, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

IP: HeLa cells,

WB: HEK-293T cells, human brain tissue, HeLa cells, U-

87 MG cells, U2OS cells

IHC: human gliomas tissue,

# **Background Information**

Reticulon (RTN) proteins are a group of membrane-bound proteins that largely reside in endoplasmic reticulum (ER). Reticulon proteins share a common sequence feature, the reticulon homology domain (RHD). Four mammalian reticulons (RTN1-4) exist. Reticulon3 (RTN3) is highly expressed in neurons. RTN3 has been shown to modulate secretory pathway protein trafficking. It is a negative modulator of BACE1 ( $\beta$ -secretase) proteolytic activity. RTN3 may induce caspase-8 cascade and apoptosis and may favor BCL2 translocation to the mitochondria upon endoplasmic reticulum stress.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Tao Wang	28985237	PLoS Pathog	WB
Sindhu Wisesa	31696206	Biochem J	IP
Zhigong Chen	32406007	J Cardiovasc Transl Res	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 12055-2-AP (RTN3 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



IP result of anti-RTN3 (IP:12055-2-AP, 4ug; Detection:12055-2-AP 1:400) with HeLa cells lysate 2000ug.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 12055-2-AP (RTN3 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 12055-2-AP (RTN3 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).