For Research Use Only

SLC25A10 Polyclonal antibody

Catalog Number:12086-1-AP

Featured Product


4 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com

	og Number: 5-1-AP	GenBank Accessi BC015797	ion Number:	Purification Method: Antigen affinity purification		
Size:		GenelD (NCBI):		Recommended Dilutions:		
550	ıg/ml	1468		WB 1:500-1:2000		
Sour		UNIPROT ID: Q9UBX3 Full Name:		IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500		
	Rabbit Isotype: IgG Immunogen Catalog Number: AG2719					
		solute carrier far	mily 25			
		(mitochondrial carrier; dicarboxylate transporter), member 10 Calculated MW: 296 aa, 32 kDa				
		Observed MW: 29-31 kDa				
ations	d Applications:		Positive Cont	ontrols:		
VVB, I	P, IHC, ELISA		WB : HEK-293 cells, A549 cells, mouse liver tissue,			
	Cited Applications: WB, IHC, IF, IP		mouse kidney tissue, HepG2 cells, PC-3 cells			
	Species Specificity:					
	IHC : human lung tissue, mouse kidney tissue Cited Species: human, rat					
TE b retri	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
catal exch level	/zes the transport of dicarboxy ange for phosphate, sulfate, an	ylates such as malat nd thiosulfate. SLC2	te and succinate acro 5A10 is expressed at	l inner membrane carrier protein whi ss the mitochondrial membrane in highest levels in kidney and liver, lo e. Two isoforms of SLC25A10 exist du		
e Publications Auth	pr P	Pubmed ID	Journal	Application		
Boro	ig Zhou 3	36104652	Hum Cell	IHC,WB		
Gaoy	uan Wang 3	32774476	Oncol Lett	WB,IHC		
Tao)	u 3	39409161	Int J Mol Sci	WB,IP,IF		


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded human lung tissue slide using 12086-1-AP (SLC 25A 10 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-SLC25A10 (IP:12086-1-AP, 3ug; Detection:12086-1-AP 1:500) with mouse liver tissue lysate 4000ug.

Various lysates were subjected to SDS PAGE followed by western blot with 12086-1-AP (SLC 25A 10 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.