For Research Use Only

## BCOR Polyclonal antibody Catalog Number: 12107-1-AP 11 Publications



| Basic Information      | Catalog Number:<br>12107-1-AP  | GenBank Accession Number<br>BC009675   | Purification Method:<br>Antigen affinity purification  |  |
|------------------------|--|--|--|--|
|                        | Concentration:<br>1100 µg/ml   | GeneID (NCBI):<br>54880  | Recommended Dilutions:<br>WB 1:2000-1:10000  |  |
|                        | Source:<br>Rabbit  | UNIPROT ID:<br>Q6W2J9  | IP 0.5-4.0 ug for 1.0-3.0 mg of total<br>protein lysate  |  |
|                        | Isotype: Full Name: IHC 1:50   IgG BCL6 co-repressor   Immunogen Catalog Number: Calculated MW:   AG2752 1703 aa, 186 kDa  | IHC 1:50-1:500   |  |  |
|                        |  |  |  |  |
|                        |  | Observed MW:<br>186 kDa  |  |  |
| Applications           | Tested Applications:   | Posit  | ive Controls:  |  |
|                        | WB, IP, IHC, ELISA<br>Cited Applications:  |  | WB : HEK-293 cells, HeLa cells, K-562 cells  |  |
|                        | WB, IHC, IP, CoIP, ChIP  |  |  |  |
|                        | Species Specificity:<br>human, mouse, rat  | IHC :  | human breast cancer tissue,  |  |
|                        | Cited Species:<br>human, mouse   |  |  |  |
|                        | Note-IHC: suggested antigen retrieval with<br>TE buffer pH 9.0; (*) Alternatively, antigen<br>retrieval may be performed with citrate<br>buffer pH 6.0   |  |  |  |
|                        | BCOR, a corepressor of BCL6, binds to transcriptional factors and specifically inhibits gene expression. BCOR plays key role in the early embryonic development, mesenchymal stem cell function and hematopiesis. BCOR can associate with HDACs, the polycomb group protein, polycomb group Ring finger protein1/nervous system polycomb1 (PCGF1/NSPC1), histone demethylase, F-box- and leucine-rich repeat protein10(FBXL10). Via repressio of TFAP2A acts as a negative regulator of osteo-dentiogenic capacity in adult stem |  |  |  |
| Background Information | polycomb1 (PCGF1/NSPC1), histo   | one demethylase, F-box- and leuc   | ine-rich repeat protein10(FBXL10). Via repressi  |  |
|                        | polycomb1 (PCGF1/NSPC1), histo   | one demethylase, F-box- and leuc   | ine-rich repeat protein10(FBXL10). Via repressi  |  |
|                        | polycomb1 (PCGF1/NSPC1), histo<br>of TFAP2A acts as a negative reg   | one demethylase, F-box- and leuc<br>ulator of osteo-dentiogenic capac  | ine-rich repeat protein10(FBXL10). Via repressi<br>ity in adult stem                             |  |
|                        | polycomb1 (PCGF1/NSPC1), hist<br>of TFAP2A acts as a negative reg<br>Author  | one demethylase, F-box- and leuc<br>ulator of osteo-dentiogenic capac<br>Pubmed ID Journal                         | ine-rich repeat protein 10(FBXL10). Via repressi<br>ity in adult stem<br>Application             |  |
| Background Information | polycomb1 (PCGF1/NSPC1), hist<br>of TFAP2A acts as a negative reg<br>Author<br>Haoqing Yang  | one demethylase, F-box- and leuc<br>ulator of osteo-dentiogenic capac<br>Pubmed ID Journal<br>31568642 Cell Prolif | ine-rich repeat protein10(FBXL10). Via repressi<br>ity in adult stem<br>Application<br>n WB,ChIP |  |


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12107-1-AP (BCOR antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 12107-1-AP (BCOR antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 12107-1-AP (BCOR antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-BCOR (IP:12107-1-AP, 4ug; Detection:12107-1-AP 1:5000) with HEK-293 cells lysate 920 ug.