For Research Use Only

SAP130 Polyclonal antibody

Catalog Number: 12130-1-AP



Basic Information

Catalog Number: GenBank Accession Number: 12130-1-AP BC017453

 Size:
 GeneID (NCBI):

 200 µg/ml
 79595

 Source:
 UNIPROT ID:

 Rabbit
 Q9H0E3

Isotype: Full Name:
IgG Sin3A-associated protein, 130kDa

Immunogen Catalog Number: Calculated MW: 110 kDa
Observed MW:

130 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200 IF 1:50-1:500

Applications

Tested Applications: IF/ICC, IHC, IP, WB, ELISA Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, HeLa cells, HepG2 cells, Jurkat

cells, K-562 cells

IP: HepG2 cells,

IHC: human pancreas cancer tissue,

IF: HepG2 cells,

Background Information

SF3b is a U2 snRNP-associated protein complex essential for spliceosome assembly. SF3b contains the spliceosomal proteins SAP 49, SAP 130, SAP 145, and SAP 155. SAP 130, SAP 145, and SAP 155 are present in a protein complex and associate with one another. While SAP 155 and SAP 130 interact with each other (directly or indirectly) within this complex, SAP 49 and SAP 145 are known to interact directly with each other. Unexpectedly, the SAP 49-SAP 145 protein-protein interaction requires the amino-terminus of SAP 49, which contains two RNA-recognition motifs. The observation that SAP 49 and SAP 145 interact directly with both U2 snRNP and the pre-mRNA suggests that this protein complex plays a role in tethering U2 snRNP to the branch site.

Storage

Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HEK-293 cells were subjected to SDS PAGE followed by western blot with 12130-1-AP (SAP130 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human pancreas cancer using 12130-1-AP (SAP130 antibody) at dilution of 1:50 (under 10x lens).



IP result of anti-SAP130 (IP:12130-1-AP, 4ug; Detection:12130-1-AP 1:600) with HepG2 cells lysate 1000ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed HepG2 cells using 12130-1-AP (SAP130 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).