## For Research Use Only

# SAP130 Polyclonal antibody

Catalog Number:12130-1-AP 1 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 12130-1-AP BC017453 GeneID (NCBI): Size: 200 ug/ml 79595 **UNIPROT ID:** Source:

Rabbit Q9H0E3 Isotype: Full Name:

Sin3A-associated protein, 130kDa

Calculated MW: Immunogen Catalog Number: AG2774 110 kDa Observed MW: 130 kDa

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB

Species Specificity: human

**Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, HepG2 cells, HeLa cells, Jurkat

**Purification Method:** 

WB 1:500-1:2000

protein lysate

IHC 1:20-1:200

IF/ICC 1:200-1:800

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

cells. K-562 cells IP: HepG2 cells,

IHC: human pancreas cancer tissue,

IF/ICC: HepG2 cells,

# **Background Information**

SF3b is a U2 snRNP-associated protein complex essential for spliceosome assembly. SF3b contains the spliceosomal proteins SAP 49, SAP 130, SAP 145, and SAP 155. SAP 130, SAP 145, and SAP 155 are present in a protein complex and associate with one another. While SAP 155 and SAP 130 interact with each other (directly or indirectly) within this complex, SAP 49 and SAP 145 are known to interact directly with each other. Unexpectedly, the SAP 49-SAP 145 protein-protein interaction requires the amino-terminus of SAP 49, which contains two RNArecognition motifs. The observation that SAP 49 and SAP 145 interact directly with both U2 snRNP and the pre-mRNA suggests that this protein complex plays a role in tethering U2 snRNP to the branch site.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Chang-Han Chen	38172660	BMC Mol Cell Biol	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




HEK-293 cells were subjected to SDS PAGE followed by western blot with 12130-1-AP (SAP130 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human pancreas cancer using 12130-1-AP (SAP130 antibody) at dilution of 1:50 (under 10x lens).



IP result of anti-SAP130 (IP:12130-1-AP, 4ug; Detection:12130-1-AP 1:600) with HepG2 cells lysate 1000ug.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SAP130 antibody (12130-1-AP) at dilution of 1:400 and CoraLite@594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4), CL488-phalloidin (green).