For Research Use Only

UBE2C Polyclonal antibody

Catalog Number: 12134-2-AP

Featured Product

16 Publications



Purification Method:

WB 1:200-1:1000

protein lysate

IHC 1:50-1:500

Positive Controls:

WB: HeLa cells,

IP: HeLa cells.

IHC: human colon cancer tissue,

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

 Catalog Number:
 GenBank Accession Number:

 12134-2-AP
 BC016292

 Size:
 GeneID (NCBI):

 1000 μ g/ml
 11065

 Source:
 UNIPROT ID:

 Rabbit
 000762

Rabbit 000762 Isotype: Full Name:

IgG ubiquitin-conjugating enzyme E2C

Immunogen Catalog Number:Calculated MW:AG2780179 aa, 20 kDaObserved MW:

20 kDa

Applications

Tested Applications: WB, IP, IHC, ELISA Cited Applications: WB, IHC, IF, CoIP Species Specificity:

human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

UBE2C(Ubiquitin-conjugating enzyme E2 C) is also known as UBCH10 and belongs to the ubiquitin-conjugating enzyme family. It is highly expressed in a variety of human primary tumors, and it is able to promote cell proliferation and malignant transformation (PMID:18703417). UBE2C, along with the E3 ligase of the anaphasepromoting complex (APC), catalyzes the ubiquitination of mitotic cyclins A and B, as well as securin. UBE2C is essential for cell cycle progression, and mutation of its active site cysteine confers a dominant-negative phenotype(PMID:17217624).

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaobo Wu	36430352	Int J Mol Sci	IHC
Zheng Cai	28560012	Am J Transl Res	WB,IHC
Jian Zhao	34249092	Front Genet	WB, IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 12134-2-AP (UBE2C antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 12134-2-AP (UBE2C Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 12134-2-AP (UBE2C Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-UBE2C (IP:12134-2-AP, 3ug; Detection:12134-2-AP 1:300) with HeLa cells lysate 2440ug.