

For Research Use Only

GAK Polyclonal antibody

Catalog Number: 12147-1-AP

1 Publications



Basic Information

Catalog Number:

12147-1-AP

Size:

600 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG2795

GenBank Accession Number:

BC008668

GeneID (NCBI):

2580

UNIPROT ID:

O14976

Full Name:

cyclin G associated kinase

Calculated MW:

134 kDa, 143 kDa

Observed MW:

150 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:250-1:1000

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : human brain tissue,

IHC : mouse brain tissue,

IF/ICC : HepG2 cells,

Background Information

GAK(cyclin-G-associated kinase) belongs to the protein kinase superfamily. GAK plays a more positive role as a modulator of signal transduction, by relaying the phosphorylation signal from an unidentified tyrosine kinase to unidentified GAK phosphorylation targets(PMID:9013862). It may be acting as an auxilin homolog that is involved in the uncoating of clathrin- coated vesicles by hsc70 in non-neuronal cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Yanchun Zhang	37196766	J Biol Chem	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 12147-1-AP (GAK antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 12147-1-AP (GAK antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 12147-1-AP (GAK antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).