For Research Use Only

Endothelin 1 Polyclonal antibody

Catalog Number:12191-1-AP

Featured Product 26 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 12191-1-AP	GenBank Accession Num BC009720	ber: Purification Method: Antigen affinity purificat	ion	
	Concentration:	GeneID (NCBI):	Recommended Dilutions:		
	650 μg/ml Source:	1906 UNIPROT ID:	WB 1:200-1:1000 IHC 1:50-1:500		
	Rabbit	P05305			
	Isotype: IgG	Full Name: endothelin 1			
	Immunogen Catalog Number: AG2847	Calculated MW: 212 aa, 24 kDa			
		Observed MW: 24 kDa			
Applications	Tested Applications:	Р	Positive Controls:		
	WB, IHC, FC (Intra), ELISA	v	WB: NIH/3T3 cells,		
	Cited Applications: WB, IHC, IF, IP	II	IHC : mouse lung tissue, mouse placenta tissue		
	Species Specificity: human, mouse				
	Cited Species: human, mouse, rat				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information		rictor peptides. Emerging basic	the endothelin/sarafotoxin family. Enc science, and animal and human data a of fibrosing disorders, including those	ll suggest f	
Notable Publications	Author	Pubmed ID Journal	Appl	ication	
	Hong Li	36180828 Mol Med	WB		
	Dandan Wang	36161694 Environ	oxicol WB		
	Peng Liu	36071548 J Cell Mo	l Med WB		


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 12191-1-AP (Endothelin 1 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse lung tissue slide using 12191-1-AP (Endothelin 1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 HUVEC cells were intracellularly stained with 0.4 ug Anti-Human Endothelin 1 (12191-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (3000-0-AP, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).