

For Research Use Only

# MVK Polyclonal antibody

Catalog Number: 12228-1-AP

7 Publications



## Basic Information

Catalog Number:

12228-1-AP

Concentration:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG2865

GenBank Accession Number:

BC016140

GeneID (NCBI):

4598

UNIPROT ID:

Q03426

Full Name:

mevalonate kinase

Calculated MW:

396 aa, 42 kDa

Observed MW:

42 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IF/ICC 1:10-1:100

## Applications

Tested Applications:

WB, IF/ICC, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, sheep

Positive Controls:

WB : Caco-2 cells, PC-3 cells, mouse uterus tissue, Jurkat cells, HepG2 cells

IF/ICC : PC-3 cells,

## Background Information

MVK (Mevalonate kinase) is a 42 kDa cytoplasmic protein that belongs to the GHMP kinase family. Mevalonate kinase catalyzes the ATP-dependent phosphorylation of mevalonic acid to form mevalonate 5-phosphate. Defects in mevalonate kinase can cause mevalonic aciduria (MEVA). It is an accumulation of mevalonic acid which causes a variety of symptoms such as psychomotor retardation, dysmorphic features, cataracts, hepatosplenomegaly, lymphadenopathy, anemia, hypotonia, myopathy and ataxia.

## Notable Publications

Author	Pubmed ID	Journal	Application
Audrey Basque	36279966	Toxicol In Vitro	WB
Min Li	32256770	Exp Ther Med	WB
Audrey Basque	35723385	Curr Issues Mol Biol	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 12228-1-AP (MVK antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of PC-3 cells using 12228-1-AP (MVK antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).



PC-3 cells were subjected to SDS PAGE followed by western blot with 12228-1-AP (MVK antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.