

For Research Use Only

CD73 Polyclonal antibody

Catalog Number: 12231-1-AP

Featured Product

59 Publications



Basic Information

Catalog Number:

12231-1-AP

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG2869

GenBank Accession Number:

BC015940

GeneID (NCBI):

4907

UNIPROT ID:

P21589

Full Name:

5'-nucleotidase, ecto (CD73)

Calculated MW:

29 kDa, 63 kDa

Observed MW:

60-70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:1000-1:3000

IHC: 1:250-1:1000

IF-P: 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF-P, ELISA

Cited Applications:

WB, IHC, IF, ELISA

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : A375 cells, human spleen tissue, K-562 cells, mouse brain tissue, HT-29 cells, HuH-7 cells, Jurkat cells

IHC : human lung cancer tissue, human stomach cancer tissue, human stomach tissue, mouse kidney tissue, human colon cancer tissue

IF-P : mouse small intestine tissue, human appendicitis tissue

Background Information

NT5E(5'-nucleotidase) is also named as NT5, NTE, CD73, ecto-5'-nucleotidase, and belongs to the 5'-nucleotidase family. It catalyzes the conversion at neutral pH of purine 5-prime mononucleotides to nucleosides, the preferred substrate being AMP. The enzyme occurs mainly as a dimer and the wide distribution of NT5E in the rat hippocampus, suggesting their involvement in the control of the purinergic signaling(PMID:17619139). This protein has two isoforms with the molecular weight of 63 kDa and 58 kDa and four glycosylation sites with a signal peptide. It can exist as a dimer(PMID:17619139). It can be detected the band of 55 kDa, 64 kDa, 70 kDa, 110 kDa by western blot(PMID:17619139; 12370277; 21533188). Defects in NT5E are the cause of calcification of joints and arteries (CALJA).

Notable Publications

| Author | Pubmed ID | Journal | Application |
|------------|-----------|---------------------|-------------|
| Xue Wu | 36330907 | Int Immunopharmacol | WB, IHC |
| Xuxin Chen | 30362554 | J Cell Physiol | |
| Wengeng Lu | 34746277 | Front Vet Sci | IF |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12231-1-AP (CD73 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 12231-1-AP (CD73 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse small intestine tissue using CD73 antibody (12231-1-AP) at dilution of 1:200 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse small intestine tissue using CD73 antibody (12231-1-AP) at dilution of 1:200 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human appendicitis tissue using CD73 antibody (12231-1-AP) at dilution of 1:200 and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).