## For Research Use Only

# NR4A1 Polyclonal antibody

Catalog Number: 12235-1-AP

**Featured Product** 

23 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 12235-1-AP BC016147
Size: GeneID (NCBI): 700 µg/ml 3164
Source: UNIPROT ID: Rabbit P22736

Rabbit P22736

Isotype: Full Name:
IgG nuclear receptor subfamily 4, group A,

Immunogen Catalog Number:

AG2875 Calculated MW: 598 aa. 64 kDa

Observed MW: 64-70 kDa

member 1

**Applications** 

Tested Applications: IF/ICC, IHC, IP, WB, ELISA Cited Applications: WB, IF, IHC, ChIP, ELISA Species Specificity:

human, mouse, rat Cited Species: human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HeLa cells, COLO 320 cells, LO2 cells, mouse liver

**Purification Method:** 

WB 1:1000-1:4000

IHC 1:500-1:2000

protein lysate

IF 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

tissue, rat liver tissue, PC-3 cells

IP: HeLa cells,

IHC: mouse colon tissue, human pancreas cancer

tissue

IF: PC-3 cells,

# **Background Information**

NR4A1 is an orphan nuclear receptor that act concomitantly with NURR1 in regulating the expression of delay-early genes during liver regeneration. Also NR4A1 has linked not only to cell apoptosis but also proliferation in response to growth factor. It plays a role in caspase-independent cell death by activation of the ERK pathway and increased activity of MEF2 transcription factors. NR4A1 exists some isoforms with MW 65 and 34 kDa.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Huimin Gao	27679973	Sci Rep	WB
Bin Huang	30224829	Sci Rep	WB
Xiaobo Wei	26497037	Mol Neurobiol	WB, IF

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 12235-1-AP (NR4A1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 12235-1-AP (NR4A1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 12235-1-AP (NR4A1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 12235-1-AP (NR4A1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 12235-1-AP (NR4A1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (10% Formaldehyde) fixed PC-3 cells using 12235-1-AP (NR4A1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-NR4A1 (IP:12235-1-AP, 4ug; Detection:12235-1-AP 1:1000) with HeLa cells lysate 1720 ug.