For Research Use Only

Cystatin C Polyclonal antibody Catalog Number: 12245-1-AP Featured Product 37 Public

37 Publications




Basic Information	Catalog Number: 12245-1-AP	GenBank Accession Number: BC013083		Purification Method: Antigen affinity purification	
	Size: GeneID (NCBI): 900 ug/ml 1471		BI):	Recommended Dilutions: WB 1:2000-1:16000	
	Source: Rabbit):	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500 IF-P 1:50-1:500 IF/ICC 1:50-1:500	al
		P01034 Full Name:			
	Isotype: IgG	cystatin C			
	Immunogen Catalog Number: AG2890	Calculated 146 aa, 16 k			
		Observed M 13 kDa	IW:		
Applications	Tested Applications: WB, IHC, IF/ICC, IF-P, IP, ELISA Cited Applications: WB, IHC, IF Species Specificity:			Positive Controls: WB : Caco-2 cells, human brain tissue, fetal human brain tissue, human milk, HepG2 cells, mouse brain tissue, HeLa cells, U-87 MG cells, U2OS cells, human urine, human serum	
			brain tissu		
			urine, hum		
	human, mouse		IP: Caco-2	IP : Caco-2 cells,	
	Cited Species: human, mouse, rat, pig, duck			IHC : human kidney tissue, human ovary tissue, mouse kidney tissue, human ovary cancer tissue	
	Note-IHC: suggested antig			IF-P : mouse kidney tissue,	
	TE buffer pH 9.0; (*) Alterr retrieval may be performe buffer pH 6.0		IF/ICC : He	eLa cells,	
Background Information	Cystatin C is a 13-kDa inhibitor of cysteine proteinases which is secreted by all cell types and is completely cleared from the organism through glomerular filtration, shown to be an early and sensitive biomarker of renal dysfunction. It is also used as an emerging biomarker in cardiovascular disease. Cystatin C is involved in a variety of inflammatory reactions. The concentration of serum cystatin C has also been shown to be unaltered in certain inflammatory conditions or other disorders of metabolism. The plasma level of serum cystatin C can be expressed as its level of generation from cells and diet and its subsequent elimination through the gut, liver, and kidneys.				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Yan Yan	26617717	Int J Clin Exp Path		
	Xiao-Lin Jiang	36309912	Aging (Albany NY) WB	
	He Tai	36279396	Aging (Albany NY) WB	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide an Aliquoting is unnecessary for -20	d 50% glycerol p	H 7.3.		


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12245-1-AP (Cystatin C antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 12245-1-AP (Cystatin C Antibody) at dilution of 1:200 (under 10x lens).




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 12245-1-AP (Cystatin C Antibody) at dilution of 1:200 (under 40x lens).




IP result of anti-Cystatin C (IP:12245-1-AP, 4ug; Detection:12245-1-AP 1:1000) with Caco-2 cells lysate 1920 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using Cystatin C antibody (12245-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).



WB result of Cystatin C antibody (12245-1-AP; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Cystatin C transfected HepG2 cells.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse kidney tissue using Cystatin C antibody (12245-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).