For Research Use Only

CCT8 Polyclonal antibody

Catalog Number:12263-1-AP

Featured Product

14 Publications




Basic Information	Catalog Number: 12263-1-AP	GenBank Accession Number: BC012584		Purification Method: Antigen affinity purification
	Size: 350 µg/ml Source: Rabbit Isotype:	GeneID (NCBI): 10694 UNIPROT ID: P50990 Full Name:		Recommended Dilutions: WB 1:2000-1:10000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IF/ICC 1:50-1:500
	IgG Immunogen Catalog Number: AG2926	chaperonin containing TCP1, subunit 8 (theta) Calculated MW: 548 aa, 60 kDa Observed MW: 60 kDa		
Applications	Tested Applications: WB, IF/ICC, IP, ELISA Cited Applications: WB, IP, IF Species Specificity: human, mouse, rat Cited Species: human, mouse, rat		Positive Controls: WB : A431 cells, mouse kidney tissue, HEK-293 cells, COLO 320 cells, HeLa cells, Jurkat cells, Raji cells IP : Jurkat cells, IF/ICC : SW480 cells, MCF-7 cells	
Background Information	CCT8, also named as C21orf112, CCTQ, KIAA0002, CCT-theta, TCP-1-theta and NY-REN-15, belongs to the TCP-1 chaperonin family. It is a molecular chaperone; assists the folding of proteins upon ATP hydrolysis. As part of the BBS/CCT complex, CCT8 may play a role in the assembly of BBSome, a complex involved in ciliogenesis regulating transports vesicles to the cilia. Known to play a role, in vitro, in the folding of actin and tubulin.(PMID:20080638)			
Notable Publications	Author	Pubmed ID	Journal	Application
	Alireza Noormohammadi	27892468	Nat Commun	WB
	S Kragness	35771867	PLoS One	WB
	Sandhya Gopal	30029015	Virology	IP
	Storage:	r after shipment.		


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12263-1-AP (CCT8 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP result of anti-CCT8 (IP:12263-1-AP, 4ug; Detection:12263-1-AP 1:500) with Jurkat cells lysate 3160 ug.



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 12263-1-AP (CCT8 Antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



WB result of CCT8 antibody (12263-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CCT8 transfected HEK-293 cells.



Immunofluorescent analysis of (4% PFA) fixed SW480 cells using CCT8 antibody (12263-1-AP) at dilution of 1:200 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594phalloidin (red).