For Research Use Only

PLXDC2 Polyclonal antibody

Catalog Number:12285-1-AP 3 Publications



Basic Information

Catalog Number: GenBank Accession Number: 12285-1-AP BC012885 GeneID (NCBI): Concentration: 700 ug/ml 84898 **UNIPROT ID:** Source: Rabbit Q6UX71 Full Name: Isotype:

plexin domain containing 2

Calculated MW: Immunogen Catalog Number: AG2940 480 aa, 54 kDa Observed MW:

60 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500

Applications

Tested Applications: WB, IHC, ELISA Cited Applications: WB, IHC, IF Species Specificity: human, mouse, rat **Cited Species:**

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, HeLa cells, MKN-45 cells

IHC: human lung cancer tissue, human liver cancer

Background Information

PLXDC2 (Plexin-domain containing 2), also named tumor endothelial marker 7-related protein (TEM7R), is a singlepass type I membrane protein with some homology to nidogen and to plexins. PLXDC2 is located in tumor endothelium and plays a role in tumor angiogenesis. Three isoforms of PLXDC2 produced by alternative splicing have been described. Recent finding revealed that PLXDC2 is a mitogen for neural progenitors and functions as a novel regulator of the development of different brain regions, suggesting its role in coordinating proliferation and differentiation in the developing nervous system.

Notable Publications

Author	Pubmed ID	Journal	Application
Yiming Guan	34124056	Front Cell Dev Biol	IHC,IF
Cheng Zhang	39197453	Dev Cell	WB
Liwei Lang	37089864	Cancer Res Commun	WB,IHC

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 12285-1-AP (PLXDC 2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 12285-1-AP (PLXDC 2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 12285-1-AP (PLXDC 2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.