For Research Use Only

Prohibitin 2 Polyclonal antibody

Catalog Number:12295-1-AP

Featured Product

52 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 12295-1-AP	GenBank Acces BC014766	ssion Number:	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI): 11331		Recommended Dilutions: WB 1:2000-1:16000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	800 ug/ml				
	Source: UNIPROTID: Rabbit Q99623				
				IHC 1:50-1:500	
	Isotype: IgG Immunogen Catalog Number: AG2977	Full Name: prohibitin 2		IF/ICC 1:400-1:1600	
		Calculated MW: 299 aa, 33 kDa			
		Observed MW: 33 kDa			
Applications	Tested Applications: WB, IHC, IF/ICC, FC (Intra), IP, ELISA		Positive Co	Positive Controls:	
			WB: HeLa c	WB : HeLa cells,	
	Cited Applications: WB, IHC, IF, IP, CoIP, Electrochemical Immunosensor Species Specificity:		IP : HeLa ce	IP : HeLa cells, mouse brain tissue	
			IHC : huma	IHC : human heart tissue, human pancreas tissue IF/ICC : HepG2 cells,	
	human, mouse	·			
	Cited Species: human, mouse, rat, rice				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	PHB2, also named as BAP, REA, D-prohibitin, BAP37 and Prohibitin-2, belongs to the prohibitin family. It acts as a mediator of transcriptional repression by nuclear hormone receptors via recruitment of histone deacetylases. PHB2 functions as an estrogen receptor (ER)-selective coregulator that potentiates the inhibitory activities of antiestrogens and represses the activity of estrogens. It competes with NCOA1 for modulation of ER transcriptional activity. PHB2 is probably involved in regulating mitochondrial respiration activity and in aging.(PMID: 10359819)				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Y Poitelon	26383514	Nat Commun	IF	
	Ryan Houston	34495738	Mol Biol Cell	WB	
	Kelly A Chamberlain	34506725	Neuron	WB	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide an		3.		


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 12295-1-AP (Prohibitin 2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




WB result of Prohibitin 2 antibody (12295-1-AP; 1:12000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Prohibitin 2 transfected HeLa cells.




IP result of anti-Prohibitin 2 (IP:12295-1-AP, 3ug; Detection:12295-1-AP 1:1000) with HeLa cells lysate 1840ug.




Immunohistochemical analysis of paraffinembedded human heart tissue slide using 12295-1-AP (Prohibitin 2 antibody at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human heart tissue slide using 12295-1-AP (Prohibitin 2 antibody at dilution of 1:200 (under 40x lens).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human Prohibitin 2 (12295-1-AP) and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Prohibitin 2 antibody (12295-1-AP) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).