## For Research Use Only

# PAX6 Polyclonal antibody

Catalog Number:12323-1-AP

**Featured Product** 

**52 Publications** 



**Basic Information** 

**Applications** 

Catalog Number: 12323-1-AP

GenBank Accession Number: BC011953

Size:

Source:

GeneID (NCBI):

5080

900 μg/ml

Full Name: paired box 6

Rabbit Isotype:

Calculated MW:

47 kDa

Immunogen Catalog Number: AG2984

Observed MW:

47 kDa

**Positive Controls:** 

**Tested Applications:** FC, IF, IHC, WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity:

human, mouse, rat

**Cited Species:** 

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

WB: Y79 cells, mouse embryo tissue, human stomach tissue, rat brain tissue

**Purification Method:** 

WB 1:500-1:3000 IHC 1:500-1:2000

Antigen affinity purification

Recommended Dilutions:

IHC: mouse brain tissue, mouse eye tissue, human retinoblastoma tissue, mouse embryo tissue

## **Background Information**

PAX6, a paired domain and homeodomain-containing transcription factor. Interaction with TRIM11 leads to ubiquitination of PAX6 and its proteasomal degradation. PAX6 is one of the earliest genes expressed in the eye field and considered a master control gene for retinal and eye development. PAX6 also regulates the development of the olfactory, central nervous systems, pituitary, and pancreas. PAX6 mutations can cause complex ocular disorders such as aniridia and Peter's anomaly.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Xin Wen	36249018	Front Oncol	WB
Xi Gu	36074953	ACS Chem Neurosci	IF
Philip G Zaworski	36058293	Anal Biochem	WB

### Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


#### Selected Validation Data




Immunofluorescent staining of PAX6 (12323-1-AP, 1:250 dilution) with 4% PFA fixed control human induced pluripotent stem cells (hiPSC) derived neuronal precursor cells (NPCs). (Green: PAX6; Blue: DAPI). Provided by BioTalentum Ltd., Hungary.




Retinal organoids (day 30) generated from human induced pluripotent stem cells (iPSCs) and fixed with 4% PFA. Stained for PAX6 with 12323-1-AP at 1:600 (green), and SOX2 with 11064-1-AP at 1:200 (red), Nuclear stain DAPI (blue). Scale bar = 20 µm. Data generated by Alessandro Bellapianta at Johannes Kepler Universitat, Austria.




Retinal organoids (day 30) generated from human induced pluripotent stem cells (iPSCs) and fixed with 4% PFA. Stained for Tubulin beta 3/TU11 using 66375-1-1g at 1:500 dilution (green) and PAX6 (12323-1-AP) at 1:500. Nuclear stain DAPI (blue). Scale bar = 50 µm. Data generated by Alessandro Bellapianta at Johannes Kepler Universitat, Austria.




Various lysates were subjected to SDS PAGE followed by western blot with 12323-1-AP (PAX6 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 12323-1-AP (PAX6 antibody) at dilution of 1:1000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 12323-1-AP (PAX6 antibody) at dilution of 1:1000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 SH-SY5Y cells were intracellularly stained with 0.4 ug Anti-Human PAX6 (12323-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).