

For Research Use Only

# VTI1A Polyclonal antibody

Catalog Number: 12354-1-AP

8 Publications



## Basic Information

Catalog Number:

12354-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG3018

GenBank Accession Number:

BC017052

GeneID (NCBI):

143187

UNIPROT ID:

Q96AJ9

Full Name:

vesicle transport through interaction with t-SNAREs homolog 1A (yeast)

Calculated MW:

217 aa, 25 kDa

Observed MW:

25-29 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IF/ICC 1:10-1:100

## Applications

Tested Applications:

WB, IF/ICC, IP, ELISA

Cited Applications:

WB, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

WB : human brain tissue, human heart tissue, human kidney tissue, human lung tissue, MCF-7 cells, mouse brain tissue

IP : MCF-7 cells,

IF/ICC : NIH/3T3 cells,

## Background Information

Fusion between membranes is mediated by specific SNARE (soluble N-ethylmaleimide-sensitive factor attachment protein receptor) complexes. Two human SNARE proteins, VTI1A and VTI1B, are homologous to the yeast Q-SNARE Vti1p which is part of several SNARE complexes in different transport steps (PMID: 12067063). VTI1A is involved in transport between the endosome and the trans-Golgi network. The SNARE complex of STX10, STX16, VTI1A, and VAMP3 is required for MPR (mannose 6-phosphate receptors) transport from endosomes to the Golgi after delivering lysosomal enzymes to the endocytic pathway (PMID: 18195106). Transport from early/recycling endosomes to the TGN is mediated by the SNARE complex of STX6, STX16, VTI1A and VAMP4 (PMID: 11839770; 21807881).

## Notable Publications

Author	Pubmed ID	Journal	Application
Yi Huo	27322682	Oncotarget	WB
Laufman Orly O	23865579	Traffic	WB,IF
Hui-Yung Song	30965672	Cells	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 12354-1-AP (VTI1A antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of NIH/3T3 cells using 12354-1-AP (VTI1A antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.



IP result of anti-VTI1A (IP:12354-1-AP, 3ug; Detection:12354-1-AP 1:1000) with MCF-7 cells lysate 520ug.