

## SF3B14 Polyclonal antibody

Catalog Number: 12379-1-AP

2 Publications

## Basic Information

## Catalog Number:

12379-1-AP

## Size:

600 µg/ml

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

AG3045

## GenBank Accession Number:

BC015463

## GeneID (NCBI):

51639

## UNIPROT ID:

Q9Y3B4

## Full Name:

splicing factor 3B, 14 kDa subunit

## Calculated MW:

14 kDa

## Observed MW:

14 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB 1:1000-1:4000

## Applications

## Tested Applications:

WB, ELISA

## Cited Applications:

WB

## Species Specificity:

human, mouse, rat

## Cited Species:

human

## Positive Controls:

WB : HeLa cells, HepG2 cells

## Background Information

Pre-mRNA splicing by the human spliceosome requires the sequential recognition by RNA and protein factors of the 5' and 3' splice sites, the branch region and the polypyrimidine tract. This sequence is characterized by the stepwise association of the U1, U2, and U4/U5/U6 snRNP particles [PMID:9476892]. Human p14 (SF3b14), a component of the spliceosomal U2 snRNP, interacts directly with the pre-mRNA branch adenosine within the context of the bulged duplex formed between the pre-mRNA branch region and U2 snRNA. This association occurs early in spliceosome assembly and persists within the fully assembled spliceosome [PMID:21062891].

## Notable Publications

Author	Pubmed ID	Journal	Application
Teng Teng	28541300	Nat Commun	WB
Frederike L Harms	37422718	Genet Med	WB

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 12379-1-AP (SF3B14 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.