### For Research Use Only

# CA8 Polyclonal antibody

Catalog Number: 12391-1-AP

**Featured Product** 

9 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 12391-1-AP BC015531

Concentration: GeneID (NCBI): 767

Source: UNIPROT ID: Rabbit P35219
Isotype: Full Name:

IgG carbonic anhydrase VIII

Immunogen Catalog Number: Calculated MW: 290 aa, 32 kDa
Observed MW:

32 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:2000-1:12000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

# **Applications**

Tested Applications:
WB, IP, IHC, ELISA
Cited Applications:
WB, IHC, IF, IP
Species Specificity:
human, mouse, rat
Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

WB: HEK-293 cells, mouse brain tissue, Jurkat cells, K-562 cells

IP: mouse brain tissue,

IHC: human gliomas tissue, mouse cerebellum tissue

# **Background Information**

The CA8 (CARP) gene encodes carbonic anhydrase VIII, which is part of a family of zinc metalloenzyme. CA8 has a central carbonic anhydrase motif, but it lacks carbonic anhydrase activity due to absence of catalytic zinc coordinating residues(PMID:2121526). CARP is a novel IP3R1-binding protein, and is expressed in Purkinje cells abundantly. CA8 is co-localized with IP3R1 in Purkinje cells and it binds to IP3R1, reducing the affinity of the receptor for its ligand, IP3(PMID:12611586). Defects in CA8 are the cause of cerebellar ataxia mental retardation and dysequilibrium syndrome type 3 (CMARQ3)(PMID:19461874).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Huai-Lu Ma	31715371	Biomed Pharmacother	WB,IHC
Ken Asada	33928345	Nucleic Acids Res	IP
Liddelow Shane A SA	23843944	PLoS One	IHC

# Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


# **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 12391-1-AP (CA8 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 12391-1-AP (CA8 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-CA8 (IP:12391-1-AP, 4ug; Detection:12391-1-AP 1:1000) with mouse brain tissue lysate 4000ug.