## For Research Use Only

# BMP4 Polyclonal antibody

Catalog Number: 12492-1-AP

**Featured Product** 

23 Publications

BC020546

GenBank Accession Number:



**Basic Information** 

Catalog Number: 12492-1-AP

Size: GeneID (NCBI):  $600 \mu \text{ g/ml}$  652

Source: UNIPROT ID: Rabbit P12644
Isotype: Full Name:

IgG bone morphogenetic protein 4

Immunogen Catalog Number: Calculated MW:

AG3191 47 kDa

Observed MW: 47 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:200-1:1000 IHC 1:50-1:500

**Applications** 

Tested Applications: WB, IHC, ELISA

Cited Applications: WB, IHC, IF

Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, goat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse heart tissue, rat heart tissue IHC: human prostate hyperplasia tissue,

# **Background Information**

BMP4, also named as BMP2B and DVR4, belongs to the TGF-beta family. BMPs have been shown to promote astroglial differentiation both in vitro and in vivo. BMP4 have been shown to be components of serum and are thought to be responsible for this differentiation. BMP4 are expressed mainly in neurons, with minimal expression detected in astrocytes and oligodendrocytes.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jia Liu	36145624	Pharmaceutics	IHC
Pengfei Xu	36435763	BMC Musculoskelet Disord	WB
Jianglan Zhang	31730904	Cancer Lett	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 12492-1-AP (BMP4 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate hyperplasia using 12492-1-AP (BMP4 antibody) at dilution of 1:50 (under 10x lens).



rat heart tissue were subjected to SDS PAGE followed by western blot with 12492-1-AP (BMP4 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.