For Research Use Only

Mu Crystallin Polyclonal antibody

Catalog Number: 12495-1-AP

3 Publications



Basic Information

Catalog Number: 12495-1-AP

Isotype:

Size: 400 µ g/ml Source: Rabbit

Immunogen Catalog Number:

AG3161

0

314 aa, 34 kDa Observed MW: 34 kDa

GenBank Accession Number:

BC018061

1428

Q14894 Full Name:

GeneID (NCBI):

UNIPROT ID:

crystallin, mu

Calculated MW:

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:4000 IHC 1:50-1:500 IF 1:50-1:500

Applications

Tested Applications: FC, IF-P, IHC, WB, ELISA

Cited Applications:

WB, IF, IHC

Species Specificity: human, mouse, rat Cited Species:

mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: Y79 cells, Jurkat cells, human kidney tissue,

human heart tissue

IHC: human gliomas tissue,

IF: human gliomas tissue,

Background Information

mu Crystallin(thiomorpholine-carboxylate dehydrogenase) is also named as THBP, CRYM, ketimine reductase and belongs to the ornithine cyclodeaminase family. This protein catalyzes the reduction of imine bonds in brain substrates that may include cystathionine ketimine (CysK) and lanthionine ketimine (LK). It is also involved in the regulation of the free intracellular concentration of triiodothyronine and access to its nuclear receptor.

Notable Publications

Author	Pubmed ID	Journal	Application
Addioi	rubilled ID	Journac	Аррисации
Yuko Ishido	26370556	Endocr J	WB
5 ··· II	2002102	N d l	WD IIIC IE
Reiji Hommyo	29603402	Neuropathology	WB,IHC,IF
John DeSisto	32634398	Dev Cell	IF

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 12495-1-AP (mu Crystallin antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human gliomas tissue using Mu Crystallin antibody (12495-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 SH-SY5Y cells were intracellularly stained with 0.2 ug Anti-Human Mu Crystallin (12495-1-AP) and Coralite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Y79 cells were subjected to SDS PAGE followed by western blot with 12495-1-AP (mu Crystallin antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.