## For Research Use Only

## SCFD1 Polyclonal antibody

Catalog Number: 12569-1-AP

**Featured Product** 

6 Publications



**Basic Information** 

Catalog Number:

12569-1-AP

Size:

500 µg/ml

Source:

Rabbit

Sotype:

GenBank Accession Number:

BC017734

GeneID (NCBI):

23256

UNIPROT ID:

Q8WVM8

Full Name:

gG sec1 family domain containing 1

Immunogen Catalog Number:Calculated MW:AG3234642 aa, 72 kDaObserved MW:

66-72 kDa

00-72 KDd

**Applications** 

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB. IP

Species Specificity: human, mouse, rat Cited Species:

human, bacteria

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: A549 cells, HEK-293T cells, human kidney tissue, human liver tissue, HEK-293 cells, HeLa cells, NIH/3T3 cells

**Purification Method:** 

WB 1:5000-1:50000

protein lysate

IHC 1:50-1:500

IF/ICC 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

IP : HeLa cells,

IHC: human endometrial cancer tissue,

IF/ICC: HeLa cells,

**Background Information** 

SCFD1, also named as C14orf163, KIAA0917, STXBP1L2, Sly1p and SLY1, belongs to the STXBP/unc-18/SEC1 family. It is a component of the syntaxin-5 SNARE complex which regulates the early secretory pathway of eukaryotic cells at the level of endoplasmic reticulum (ER) to Golgi transport. (PMID:21242315) SCFD1 plays a role in SNARE-pin assembly and Golgi-to-ER retrograde transport via its interaction with COG4. It is involved in vesicular transport between the endoplasmic reticulum and the Golgi.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Xue Shen	30408277	J Appl Microbiol	WB
Guizhen Wang	32687939	Microb Pathog	WB
Hong Huang	35465820	Autophagy	WB,IP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.cor

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 12569-1-AP (SCFD1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




IP result of anti-SCFD1 (IP:12569-1-AP, 3ug; Detection:12569-1-AP 1:500) with HeLa cells lysate 2150ug.



Immunohistochemical analysis of paraffinembedded human endometrial cancer tissue slide using 12569-1-AP (SCFD1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human endometrial cancer tissue slide using 12569-1-AP (SCFD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using 12569-1-AP (SCFD1 antibody) at dilution of 1:50 and Coralite488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).