For Research Use Only

GCLC Polyclonal antibody

Catalog Number: 12601-1-AP

Featured Product

131 Publications



Basic Information

Catalog Number:

12601-1-AP

BC022487

Concentration:

800 ug/ml

2729

Source:

UNIPROT ID:

Rabbit

P48506

Isotype:

GenBank Accession Number:

BC022487

GeneID (NCBI):

2729

UNIPROT ID:

Full Name:

glutamate-cysteine ligase, catalytic

Immunogen Catalog Number: subunit

AG3252 Calculated MW: 637 aa, 73 kDa

Observed MW: 73 kDa

Applications

Tested Applications:
WB, IHC, IP, ELISA
Cited Applications:
WB, IHC, IF, IP
Species Specificity:
human, mouse, rat
Cited Species:
human, mouse, rat, goat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Antigen affinity purification
Recommended Dilutions:
WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

Purification Method:

Positive Controls:

WB: Daudi cells, HEK-293 cells, LO2 cells, mouse kidney tissue, Jurkat cells, Raji cells, mouse lung tissue, rat lung tissue

IP: mouse kidney tissue,

IHC: human kidney tissue, human liver tissue

Background Information

GCLC (Glutamate--cysteine ligase catalytic subunit) belongs to the glutamate--cysteine ligase type 3 family. It is also named as GLCL, GLCLC. It is the first rate-limiting enzyme in glutathione (GSH) biosynthesis. GCLC) gene may be associated with coronary endothelial vasomotor dysfunction and myocardial infarction (MI) (PMID:12598062). Defects in GCLC are the cause of hemolytic anemia (HAGGSD).

Notable Publications

Author	Pubmed ID	Journal	Application
Lu Wang	34555641	Int Immunopharmacol	WB
Ana Tomasovic	27645114	Matrix Biol	WB
Juan Chen	34577027	Molecules	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12601-1-AP (GCLC antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 12601-1-AP (GCLC Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 12601-1-AP (GCLC Antibody) at dilution of 1:200 (under 40x lens).



IP result of anti-GCLC (IP:12601-1-AP, 4ug; Detection:12601-1-AP 1:5000) with mouse kidney tissue lysate 2160 ug.