For Research Use Only

golgin 97 Polyclonal antibody Catalog Number:12640-1-AP 27 Publications




Basic Information	Catalog Number: 12640-1-AP	GenBank Accession Number: BC032853		Purification Method:					
	Size:	GenelD (NC	BI).	Antigen affinity purification Recommended Dilutions: WB 1:2000-1:16000 IF/ICC 1:500-1:2000					
	800 µ g/ml	2800	<i>bij</i> .						
	Source:	UNIPROT ID):						
	Rabbit	Q92805							
	Isotype:	Full Name: golgi autoantigen, golgin subfamily a, 1 Calculated MW: 767 aa, 88 kDa Observed MW: 97 kDa							
	IgG Immunogen Catalog Number: AG3335								
					Applications	Tested Applications:		Positive Co	ntrols:
						WB IE/ICC EC (Intra) ELISA			93 cells, SMMC-7721 cells, HeLa cells,
	Cited Applications:						HepG2 cells		
WB, IHC, IF, IP IF/ICC : Hel			.a cells,						
Species Specificity: human Cited Species: human, mouse, monkey									
				Background Information		Golgin subfamily A member 1 is to interact with ARL1. GOLGA1 is Catalog#12640-1-AP specially r	associated with S	Sjogren's syndrome. The	he GOLGA1 gene. GOLGA1 has been show e MW of this protein is 97 kDa, and nows strong Gogli staining.
Notable Publications	Author	Pubmed ID	Journal	Application					
	Mengyue You	36252341	Redox Biol	IF					
	Chengfeng Merriman	31591269	J Biol Chem	WB					
	Rachel E Turn	34818063	Mol Biol Cell	IHC					
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide ar Aliquoting is unnecessary for -20	nd 50% glycerol p							


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 12640-1-AP (golgin 97 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using golgin 97 antibody (12640-1-AP) at dilution of 1:1000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



1X10^{^6} HeLa cells were intracellularly stained with 0.4 ug Anti-Human golgin 97 (12640-1-AP) and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).