For Research Use Only

SNAP29 Polyclonal antibody

Catalog Number: 12704-1-AP

Featured Product

30 Publications



Basic Information

Catalog Number: GenBank Accession Number: 12704-1-AP BC009715

Concentration: GeneID (NCBI): 9342

Source: UNIPROT ID:

Rabbit 095721

Isotype: Full Name: synaptosomal-associated protein,

Immunogen Catalog Number: 29kDa

AG3382 Calculated MW:

258 aa, 29 kDa Observed MW: 29 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF/ICC 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity: human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: human kidney tissue, K-562 cells, human liver tissue, Jurkat cells, HEK-293 cells

IP: Jurkat cells.

IHC: mouse kidney tissue, human kidney tissue,

human stomach tissue

IF/ICC: PC-12 cells, HEK-293 cells, HeLa cells

Background Information

SNAREs, soluble N-ethylmaleimide-sensitive factor-attachment protein receptors, are essential proteins for the fusion of cellular membranes. SNAREs localized on opposing membranes assemble to form a trans-SNARE complex, an extended, parallel four alpha-helical bundle that drives membrane fusion. SNAP29 is a SNARE involved in autophagy through the direct control of autophagosome membrane fusion with the lysosome membrane. SNAP29 also plays a role in ciliogenesis by regulating membrane fusions.

Notable Publications

Author	Pubmed ID	Journal	Application
Zheng Li	31775910	Stem Cell Res Ther	WB
Hao-Long Zeng	29174846	Biochim Biophys Acta	WB
Yonglu Che	31712554	Nat Commun	WB,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




human kidney tissue were subjected to SDS PAGE followed by western blot with 12704-1-AP (SNAP29 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (-20°C Ethanol) fixed PC-12 cells using SNAP29 antibody (12704-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 12704-1-AP (SNAP29 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 12704-1-AP (SNAP29 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-SNAP29 (IP:12704-1-AP, 4ug; Detection:12704-1-AP 1:600) with Jurkat cells lysate 3200ug.



Immunohistochemical analysis of paraffinembedded normal stomach slide using 12704-1-AP (SNAP29 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).