For Research Use Only

BBS4 Polyclonal antibody

Catalog Number: 12766-1-AP

Featured Product

20 Publications

BC027624

UNIPROT ID:

Q96RK4 Full Name:

GenBank Accession Number:



Basic Information

Catalog Number: 12766-1-AP

Size: GeneID (NCBI): 550 ug/ml 585

Source: Rabbit Isotype:

gG Bardet-Biedl syndrome 4

Immunogen Catalog Number:Calculated MW:AG3430519 aa, 58 kDaObserved MW:

50-60 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:4000 IHC 1:20-1:200

IF/ICC 1:50-1:400

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IF

Species Specificity: human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: HEK-293 cells, mouse kidney tissue

IHC: human kidney tissue,

IF/ICC: hTERT-RPE1 cells,

Background Information

BBS4 (Bardet-Biedl syndrome protein 4) is one of the early BBS proteins discovered, with 35 mutations reported to cause the BBS phenotype (PMID: 26762677). BBS4 undergoes localization to the centriolar satellites of centrosomes and basal body of primary cilia, where it takes part in recruiting cargo to centriolar satellites and allowing the formation of a functional centrosomal microtubule organizing center (MTOC) (PMID: 25860617). Some research has found that BBS4 expression in humans during the development of CNS and sensory organs, concerning the cardinal clinical signs of BBS (PMID: 33860840).

Notable Publications

Author	Pubmed ID	Journal	Application
Naila Haq	31479441	PLoS Biol	WB,IF
Deniz Conkar	31582766	Sci Rep	WB
Tianyu Wu	36395215	Science	IF

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HEK-293 cell lysates were subjected to SDS PAGE followed by western blot with 12766-1-AP (BBS4 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




IF result (cytoplasm and the length of cilia cilia stain) of anti-BBS4 (12766-1-AP; 1:50) with serum-starved hTERT-RPE1 cell (MeOH fixed) by Dr. Moshe



Immunohistochemical analysis of paraffinembedded human kidney using 12766-1-AP (BBS4 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney using 12766-1-AP (BBS4 antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (4% PFA) fixed hTERT-RPE1 cells using BBS4 antibody (12766-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CoraLite® 594 ARL13B antibody (CL594-17711, red).