# For Research Use Only

# MAFF Polyclonal antibody

Catalog Number: 12771-1-AP

**Featured Product** 

**14 Publications** 



**Basic Information** 

Catalog Number:

12771-1-AP

Size:

800 µg/ml

23764

Source:

Rabbit

Q9ULX9

Isotype:

GenBank Accession Number:

BC015037

GeneID (NCBI):

23764

UNIPROT ID:

Q9ULX9

Full Name:

IgG v-maf musculoaponeurotic
Immunogen Catalog Number: fibrosarcoma oncogene homolog F

AG3460 (avian)

Calculated MW: 17 aa, 8 kDa Observed MW: 18-23 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2400

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:20-1:200

**Applications** 

Tested Applications:
WB, IP, IHC, ELISA
Cited Applications:
WB, IHC, IF, CoIP
Species Specificity:
human, mouse, rat
Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: human lung tissue, IP: SKOV-3 cells,

IHC: human ovary tumor tissue, human lung cancer

tissue

# **Background Information**

MAFF encodes a basic region and leucine zipper (bZIP)-type transcription factor protein (PMID: 8361754). MAFF interacts with the upstream promoter region of oxytocin receptor (OXTR) and plays a role in OXTR upregulation (PubMed: 7798245, PubMed:10527846). Mice lacking MAFF, MAFG and MAFK are embryonic lethal (PMID: 22158967). MAFF has a higher expression in adipose, colon, lung, prostate and skeletal muscle tissues (PMID: 24304889).

#### **Notable Publications**

| Author         | Pubmed ID | Journal                    | Application |
|----------------|-----------|----------------------------|-------------|
| Masahiro Okabe | 30379099  | Am J Physiol Renal Physiol | IF          |
| Tianwei Chen   | 36453019  | Mol Oncol                  | WB          |
| Shuyuan Shen   | 32322670  | Mol Ther Oncolytics        | WB          |

### Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


# **Selected Validation Data**




IP result of anti-MAFF (IP:12771-1-AP, 4ug; Detection:12771-1-AP 1:500) with SKOV-3 cells lysate 3200 ug.




human lung tissue were subjected to SDS PAGE followed by western blot with 12771-1-AP (MAFF antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor using 12771-1-AP (MAFF antibody) at dilution of 1:100 (under 10x lens)



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 12771-1-AP (MAFF antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 12771-1-AP (MAFF antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).