For Research Use Only

LRAT Polyclonal antibody Catalog Number:12815-1-AP 7 Publications




Basic Information	Catalog Number: 12815-1-AP	GenBank Accession Number: BC031053	Purification Method: Antigen affinity purification	
	Concentration:	GenelD (NCBI):	Recommended Dilutions:	
	500 µg/ml	9227	WB 1:500-1:1000	
	Source: Rabbit	UNIPROT ID: 095237		
	Isotype:	Full Name:		
	IgG Immunogen Catalog Number: AG3556	lecithin retinol acyltransferase (phosphatidylcholineretinol O- acyltransferase)		
		Calculated MW: 230 aa, 26 kDa		
		Observed MW:		
		26 kDa, 50-54 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, ELISA	WB : mouse testis tissue, Y79 cells		
	Cited Applications: WB			
	Species Specificity: human, mouse, rat			
	Cited Species:			
	human, mouse, rat			
Background Information	LRAT belongs to the H-rev107 family. LRAT transfers the acyl group from the sn-1 position of phosphatidylcholine all-trans retinol, producing all-trans retinyl esters. LRAT plays a critical role in vision. It provides the all-trans retin ester substrates for the isomerohydrolase which processes the esters into 11-cis-retinol in the retinal pigment epithelium; due to a membrane-associated alcohol dehydrogenase, LRAT monomers (26kDa) interact in membrar and form functional homodimers (50-54 kDa) through protein-protein interactions and disulfide bond formation.Defects in LRAT are a cause of severe early-onset retinal dystrophy (RD). This antibody was generated against the internal residues of human LRAT. It is predicted to detect endogenous LRAT.			
	Author	Pubmed ID Journal	Application	
Notable Publications				
Notable Publications		J Food Biochem	WB	
Notable Publications	Wei Li	31353518 J Food Biochem 28734946 Am J Pathol	WB WB	
Notable Publications	Wei Li Volha V Malechka			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse testis tissue were subjected to SDS PAGE followed by western blot with 12815-1-AP (LRAT Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.