For Research Use Only

NUDT16 Polyclonal antibody

Catalog Number: 12889-1-AP

Featured Product

6 Publications



Purification Method:

WB 1:1000-1:6000

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number:

12889-1-AP

Size:

GeneID (NCBI):

1100 ug/ml

131870

Source:

Rabbit

Q96DE0

Isotype:

GeneID (NCBI):

171870

GeneID (NCBI):

171870

Full Name:

Full Name: IF/ICC 1:200-1:800 nudix (nucleoside diphosphate linked

Immunogen Catalog Number: moiety X)-type motif 16

AG3925 Calculated MW:

195 aa, 21 kDa Observed MW: 21 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB, IHC

IgG

Species Specificity: human

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HeLa cells, K-562 cells

IP: HEK-293 cells,

IHC: human kidney tissue, IF/ICC: HEK-293 cells,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Lu Wang	36389726	Front Immunol	IHC
Christopher R Neil	34613784	Mol Biol Cell	WB
Siddharth Shukla	25071210	Proc Natl Acad Sci U S A	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12889-1-AP (NUDT16 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




IP result of anti-NUDT16 (IP:12889-1-AP, 3ug; Detection:12889-1-AP 1:1000) with HEK-293 cells lysate 2400ug.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 12889-1-AP (NUDT16 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 12889-1-AP (NUDT16 Antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using NUDT16 antibody (12889-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).