For Research Use Only

CUL1 Polyclonal antibody Catalog Number:12895-1-AP Featured Product

Featured Product






Basic Information	Catalog Number: GenBank Accession Number: 12895-1-AP BC034318		sion Number:	Purification Method: Antigen affinity purification		
	Size:	GeneID (NCBI):		Recommended Dilutions:		
	350 µg/ml	8454	•54		WB 1:5000-1:50000 IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	Source: UNIPROT ID: Rabbit Q13616 Isotype: Full Name:			protein lysate IHC 1:250-1:1000		
	lgG	cullin 1				
	Immunogen Catalog Number: AG3585	Calculated MW 90 kDa	Calculated MW: 90 kDa Observed MW: 80-90 kDa			
Applications	Tested Applications:		Positive Controls:			
	WB, IHC, FC (Intra), IP, ELISA		WB : A549 cells, BxPC-3 cells, K-562 cells			
	Cited Applications: IP : K-562 WB, IP, CoIP		IP : K-562 cel	cells, HeLa cells		
	Species Specificity: human, mouse		IHC : human lung cancer tissue,			
	Cited Species: human, mouse					
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	The cullin family proteins are scaffold proteins for the Ring finger type E3 ligases. Humans express seven cullin proeins: CUL1-3, CUL4A, CUL4B, CUL5, and CUL7. Each cullin protein can form an E3 ligase similar to the prototype Ring-type E3 ligase Skp1-CUL1-F-box complex. The Cullin-RING-finger type E3 ligases are important regulators in early embryonic development, as highlighted by genetic studies demonstrating that knock-out of CUL1, CUL3, or CUL4A in mice results in early embryonic lethality.					
Notable Publications	Author	Pubmed ID	Journal		Application	
	Tiantian Xu	36253371	Signal Transduct Tai	rget Ther	WB	
	Lu Yin	36240645	Neoplasia		WB	
	Tomoaki Nagai	30404837	J Cell Sci		WB	


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 12895-1-AP (CUL1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 12895-1-AP (CUL1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 12895-1-AP (CUL1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-CUL1 (IP:12895-1-AP, 4ug; Detection:12895-1-AP 1:2000) with K-562 cells lysate 1760 ug.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human CUL1 (12895-1-AP) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).