For Research Use Only

CYP2W1 Polyclonal antibody

Catalog Number: 12904-1-AP

Featured Product

1 Publications



Basic Information

Catalog Number: GenBank Accession Number: 12904-1-AP BC025761 GeneID (NCBI): Size: 400 ug/ml 54905 **UNIPROT ID:** Source: Rabbit

Q8TAV3 Full Name:

cytochrome P450, family 2, subfamily W, polypeptide 1

AG3803 Calculated MW: 54 kDa

Observed MW: 45-54 kDa

Applications

Tested Applications: WB, IHC, ELISA **Cited Applications:**

Immunogen Catalog Number:

Isotype:

Species Specificity: human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000 IHC 1:50-1:500

Positive Controls:

WB: HEK-293 cells, HepG2 cells, NIH/3T3 cells IHC: human stomach cancer tissue, human colon

cancer tissue

Background Information

Cytochrome P450s have been proposed to be of importance both in carcinogenesis, by activating precarcinogens, and as determinants of cancer chemotherapy, where they participate in activation or inactivation of anti-cancer drugs. CYP2W1 is exclusively expressed in transformed tissue in the adult human, mainly in colon tumors. It is a potential drug target in cancer therapy. CYP2W1 has a molecular weight of 52-54 kDa, and 30/45 kDa of potential isoforms.

Notable Publications

Author	Pubmed ID	Journal	Application
Hanna Szaefer	31561186	Pharmacol Rep	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




WB result of CYP2W1 antibody (12904-1-AP; 1:300; incubated at room temperature for 1.5 hours) with sh-Control and sh-CYP2W1 transfected HEK-293 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 12904-1-AP (CYP2W1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded stomach cancer slide using 12904-1-AP (CYP2W1 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded Human Colon cancer tissue slide using 12904-1-AP (CYP2W1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).