

For Research Use Only

HDAC2 Polyclonal antibody

Catalog Number: 12922-3-AP

Featured Product

87 Publications



Basic Information

Catalog Number:

12922-3-AP

Concentration:

600 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG3607

GenBank Accession Number:

BC031055

GeneID (NCBI):

3066

UNIPROT ID:

Q92769

Full Name:

histone deacetylase 2

Calculated MW:

458 aa, 52 kDa; 488 aa, 55 kDa

Observed MW:

55-60 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:2000-1:8000

IF/ICC 1:1000-1:4000

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, ChIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Positive Controls:

WB : HEK-293 cells, human kidney tissue, MCF-7 cells, rat liver tissue, HeLa cells, HepG2 cells, L02 cells, C6 cells, NIH/3T3 cells, rat kidney tissue

IP : HEK-293 cells,

IHC : mouse brain tissue,

IF/ICC : HepG2 cells, HeLa cells

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Histone deacetylases (HDAC) are a class of enzymes that remove the acetyl groups from the lysine residues leading to the formation of a condensed and transcriptionally silenced chromatin. Histone deacetylases act via the formation of large multiprotein complexes, and are responsible for the deacetylation of lysine residues at the N-terminal regions of core histones (H2A, H2B, H3 and H4). At least 4 classes of HDAC were identified. As a class I HDAC, HDAC2 was primarily found in the nucleus. HDAC2 forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus, it plays an important role in transcriptional regulation, cell cycle progression and developmental events. This antibody is a rabbit polyclonal antibody raised against residues near the C terminus of human HDAC2.

Notable Publications

Author	Pubmed ID	Journal	Application
Hong Mai	34586697	J Cell Mol Med	IHC
Daniel B McClatchy	32994440	Sci Rep	WB
Z Li	26411366	Oncogene	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12922-3-AP (HDAC2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




IP result of anti-HDAC2 (IP:12922-3-AP, 4ug; Detection:12922-3-AP 1:20000) with HEK-293 cells lysate 960 ug.




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using HDAC2 antibody (12922-3-AP) at dilution of 1:2000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).




1x10⁶ HeLa cells were intracellularly stained with 0.25 ug HDAC2 Polyclonal antibody (12922-3-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 12922-3-AP (HDAC2 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 12922-3-AP (HDAC2 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HDAC2 antibody (12922-3-AP) at dilution of 1:1000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).