### For Research Use Only

# NFYA Polyclonal antibody

Catalog Number: 12981-1-AP

**Featured Product** 

4 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 12981-1-AP BC039244 GeneID (NCBI): Size: 800  $\mu$  g/ml 4800 **UNIPROT ID:** Source: Rabbit P23511 Full Name: Isotype:

nuclear transcription factor Y, alpha

Calculated MW: Immunogen Catalog Number:

AG3604 318aa,34 kDa; 347aa,39 kDa

> Observed MW: 40 kDa, 46 kDa

**Applications** 

**Tested Applications:** IF/ICC, IHC, WB, ELISA Cited Applications: IHC. WB

Species Specificity: human, mouse **Cited Species:** human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, HeLa cells, HT-1080 cells, Jurkat

**Purification Method:** 

WB 1:500-1:1000 IHC 1:250-1:1000

IF 1:10-1:100

Antigen affinity purification

Recommended Dilutions:

cells. K-562 cells

IHC: human colon cancer tissue, human prostate

cancer tissue IF: HepG2 cells,

# **Background Information**

NFYA, also named as Nuclear transcription factor Y subunit alpha, is a 347 amino acid protein, which contains 1 NFYA/HAP2-type DNA-binding domain and belongs to the NFYA/HAP2 subunit family. NFYA exists the two 41- to 45-kDa isoforms of NF-YA ( 11287613). NFYA is a component of the sequence-specific heterotrimeric transcription factor (NF-Y) which specifically recognizes a 5'-CCAAT-3' box motif found in the promoters of its target genes. NF-Y can function as both an activator and a repressor, depending on its interacting cofactors. NF-YA positively regulates the transcription of the core clock component ARNTL/BMAL1

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Joshua E Burda	35614216	Nature	IHC
Bin Cao	30106137	Int J Oncol	WB
Matthew S Fuller	28446681	J Virol	WB

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**

 $116kd \rightarrow \\ 66kd \rightarrow \\ 45kd \rightarrow \\ \\ 35kd \rightarrow \\ \\$ 

HEK-293 cells were subjected to SDS PAGE followed by western blot with 12981-1-AP (NFYA antibody) at dilution of 1:200 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells, using NFYA antibody 12981-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 12981-1-AP (NFYA antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).