

For Research Use Only

PHD1 Polyclonal antibody

Catalog Number: 12984-1-AP

11 Publications



Basic Information

Catalog Number:

12984-1-AP

Concentration:

600 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG3616

GenBank Accession Number:

BC036051

GeneID (NCBI):

112398

UNIPROT ID:

Q96K50

Full Name:

egl nine homolog 2 (C. elegans)

Calculated MW:

407 aa, 44 kDa

Observed MW:

48 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse

Cited Species:

human, mouse, rat

Positive Controls:

WB : mouse testis tissue, HEK-293 cells, HeLa cells, human spleen tissue, mouse heart tissue, PC-3 cells

Background Information

EGLN2, also named as PHD1, EIT6 and HPH-3, catalyzes the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. It hydroxylates HIF-1 alpha at 'Pro-402' and 'Pro-564', and HIF-2 alpha. EGLN2 functions as a cellular oxygen sensor and, under normoxic conditions, targets HIF through the hydroxylation for proteasomal degradation via the von Hippel-Lindau ubiquitination complex. It may play a role in cell growth regulation.

Notable Publications

Author	Pubmed ID	Journal	Application
Suping Li	34687132	J Cell Mol Med	IF
Haixu Xu	35589690	Cell Death Dis	WB
Changhui Yu	35351157	Respir Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse testis tissue were subjected to SDS PAGE followed by western blot with 12984-1-AP (PHD1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.