For Research Use Only

TFAP2A/AP-2 Polyclonal antibody

Catalog Number:13019-3-AP

Featured Product

10 Publications



Purification Method:

WB 1:2000-1:10000 IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: 13019-3-AP Size: 450 ug/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG4112

GenBank Accession Number:

BC017754

GeneID (NCBI):
7020

UNIPROT ID:
P05549

transcription factor AP-2 alpha (activating enhancer binding protein

2 alpha)

Full Name:

Calculated MW: 431 aa, 47 kDa Observed MW: 47 kDa

Positive Controls:

WB: HepG2 cells, Y79 cells, MCF-7 cells

IF/ICC: HepG2 cells,

Applications

Tested Applications: WB, IF/ICC, ELISA Cited Applications: WB, IP, ChIP Species Specificity:

human
Cited Species:
human, mouse

Background Information

The activator protein-2 (AP-2) family of transcription factors comprises five 52-kDa isoforms (AP-2 α , AP-2 β , AP-2 γ , AP-2 δ , and AP-2 ϵ), which share a common structure: a proline/glutamine-rich transactivation domain in the N-terminal region and a helix-span-helix domain in the C-terminal region, which mediates dimerization and site-specific DNA binding. Depending on the cellular context, the AP-2 transcription factors are individually associated either with cell differentiation and development or with cancer progression/regression (PMID: 21966377). TFAP2A (AP-2-alpha) is the only AP-2 protein required for early morphogenesis of the lens vesicle. Together with the CITED2 coactivator, stimulates the PITX2 P1 promoter transcription activation (PMID: 11694877).

Notable Publications

Author	Pubmed ID	Journal	Application
Ming-Xu Qiao	27499261	Oncol Rep	
Zhan-Ping Lu	26073327	Acta Pharmacol Sin	WB, ChIP
Yanlu Xiong	33824285	Cell Death Dis	ChIP

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 13019-3-AP (TFAP2A/AP-2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using TFAP2A/AP-2 antibody (13019-3-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).