## For Research Use Only

# RAD9A Polyclonal antibody

Catalog Number: 13035-1-AP 5 Publications



**Purification Method:** 

WB 1:1000-1:4000 IHC 1:400-1:1600

IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number:

13035-1-AP

BC014848

Size:

GeneID (NCBI):

450 µg/ml

5883

Source:

UNIPROT ID:

Rabbit

Q99638

Isotype:

GenBank Accession Number:

BC014848

UNCBI):

450 µg/ml

5883

Full Name:

gG RAD9 homolog A (S. pombe)

Immunogen Catalog Number: Calculated MW: AG3687 391 aa, 42 kDa Observed MW:

55-60 kDa

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications:

WB, IHC

Species Specificity:

human
Cited Species:
human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, A549 cells, MCF-7 cells IHC: human breast cancer tissue,

IF/ICC: HeLa cells,

# **Background Information**

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Xiaoli Xu	30184152	J Mol Cell Biol	WB
Xu Zhao	34042508	DNA Cell Biol	WB
Bin Chen	35152263	Oncogene	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




HeLa cells were subjected to SDS PAGE followed by western blot with 13035-1-AP (RAD9A antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 13035-1-AP (RAD9A antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 13035-1-AP (RAD9A antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 13035-1-AP (RAD9A antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).