

For Research Use Only

MCM4 Polyclonal antibody

Catalog Number: 13043-1-AP

15 Publications



Basic Information

Catalog Number:

13043-1-AP

Size:

350 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG3703

GenBank Accession Number:

BC031061

GeneID (NCBI):

4173

UNIPROT ID:

P33991

Full Name:

minichromosome maintenance
complex component 4

Calculated MW:

863 aa, 97 kDa

Observed MW:

97 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 µg for 1.0-3.0 mg of total
protein lysate

IHC 1:1000-1:4000

Applications

Tested Applications:

WB, IP, IHC, ELISA

Cited Applications:

WB

Species Specificity:

human

Cited Species:

human, mouse

Positive Controls:

WB : HeLa cells, HEK-293 cells, HL-60 cells, human liver
tissue

IP : HEK-293 cells,

IHC : human colon cancer tissue, mouse testis tissue,
rat colon tissue

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Background Information

DNA replication licensing factor MCM4 (MCM4), also named Cdc21, acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity. This antibody is a rabbit polyclonal antibody raised against the C-terminal 350 aa sequence of MCM4 protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Nathalie Eisenhardt	26524493	Nat Struct Mol Biol	WB
Hongyuan Song	29908342	Biomaterials	WB
Lu Yan	31413757	J Cancer	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 13043-1-AP (MCM4 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




IP result of anti-MCM4 (IP:13043-1-AP, 4ug; Detection:13043-1-AP 1:600) with HEK-293 cells lysate 1200ug.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 13043-1-AP (MCM4 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 13043-1-AP (MCM4 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 13043-1-AP (MCM4 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 13043-1-AP (MCM4 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat colon tissue slide using 13043-1-AP (MCM4 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat colon tissue slide using 13043-1-AP (MCM4 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).